

Long-Acting Lenacapavir in People With Multidrug-Resistant HIV-1: Week 52 Results




Ogbuagu, Onyema¹, Sorana Segal- Maurer², Cynthia Brinson³, Ploenchan Chetchotisakd⁴, Joseph P. McGowan⁵, Kimberly Workowski⁶, Hui Wang⁷, Nicolas Margot⁷, Hadas Dvory-Sobol⁷, Martin S Rhe⁷, Jared Baeten⁷, Lindsey Griffiths⁸, Jean-Michel Molina⁹

¹Yale University, School of Medicine, New Haven, CT, US; ²New York Presbyterian Queens, Flushing, NY, US; ³Central Texas Clinical Research, Austin, TX, US; ⁴Srinagarind Hospital, Khon Kaen, Thailand; ⁵North Shore University Hospital, Manhasset, NY, US; ⁶Emory University, Atlanta, GA; ⁷Gilead Sciences, Inc., Foster City, CA; ⁸Gilead Sciences, Pty Ltd, Melbourne AUS; ⁹Hopital Saint Louis, Paris, France

Introduction Results

Lenacapavir (LEN; GS-6207) Targets Multiple Stages of HIV Replication Cycle^{1,2}




- LEN is a novel, highly potent, long-acting, first-in-class, HIV-1 capsid inhibitor
- LEN can meet significant unmet medical needs:
 - A new mechanism of action for heavily treatment-experienced (HTE) people with multidrug-resistant (MDR) HIV-1 and limited treatment options
 - Reduction of daily pill burden through less frequent dosing for treatment and prevention
- Highly desirable *in vitro* profile with picomolar antiviral activity (EC₅₀: 50–100 pM)
 - Retains full activity against nucleoside reverse-transcriptase inhibitor (NRTI)-, non-NRTI (NNRTI)-, integrase strand transfer inhibitor (INSTI)-, and protease inhibitor (PI)-resistant mutants³⁻⁵
 - No observed pre-existing resistance⁶
- In treatment-naïve people with HIV-1, LEN + emtricitabine/tenofovir alafenamide led to 94% virologic suppression at Week 28⁷
- Previously in the CAPELLA Study (NCT04150068) in HTE people with MDR HIV-1:
 - LEN achieved its primary endpoint as a functional monotherapy when added to a failing regimen⁸:
 - Participants with ≥0.5-log₁₀ decline: LEN 88% vs placebo 17% (p<0.001)
 - Mean HIV-1 RNA decline: LEN 1.9 vs placebo 0.3 log₁₀ (p<0.001)
 - LEN + optimised background regimen (OBR) led to 81% virologic suppression at Week 26⁹

Objectives

- To evaluate the safety and efficacy (using the FDA Snapshot algorithm) of LEN in combination with an OBR at Weeks 26 and 52

Methods



- Week 52 efficacy was summarised only for the randomised cohort (n=36), as most participants in the non-randomised cohort have not yet reached Week 52
- Safety was summarised for both the randomised and non-randomised cohorts (N=72)

Methods


Baseline Characteristics

	Randomised		Non-randomised	Total: N=72
	LEN: n=24	Placebo: n=12	LEN: n=36	
Age, median (range), years	55 (24–71)	54 (27–59)	49 (23–78)	52 (23–78)
Sex, % female at birth	29	25	22	25
Race, % Black	42	55	31	38
Ethnicity, % Hispanic/Latinx	25	36	14	21
HIV-1 RNA, median (range), log ₁₀ c/mL	4.2 (2.3–5.4)	4.9 (4.3–5.3)	4.5 (1.3–5.7)	4.5 (1.3–5.7)
>75,000 c/mL, %	17	50	28	28
CD4 count, median (range), cells/μL	172 (16–827)	85 (6–237)	195 (3–1296)	150 (3–1296)
<200 cells/μL, %	67	92	53	64
No. of prior ARV agents, median (range)	9 (2–24)	9 (3–22)	13 (3–25)	11 (2–25)
No. of fully active agents in OBR, %				
0	17	17	17	17
1	29	58	36	38
≥2	54	25	47	46
Known resistance to ≥2 drugs in class, %				
NRTI	96	100	100	99
NNRTI	92	100	100	97
INSTI	83	58	64	69
PI	83	67	83	81


CD4, cluster of differentiation-4.

Results


Efficacy at Week 26 in Randomised and Non-randomised Cohorts



Efficacy in Randomised Cohort (n=36)



Efficacy by No. of Fully Active Agents in OBR at Week 52 in Randomised Cohort (n=36)




Efficacy by No. of Fully Active Agents in OBR at Week 52 in Randomised Cohort (n=36)

n (%)	Randomised Cohort: n=36 (presented at IAS 2021, EACS 2021) ^{9,10}	Non-randomised Cohort: n=36
Participants meeting criteria for resistance testing	11 (31)	10 (28)
Emergent LEN resistance [†]	4 (11)	4 (11)
M66I	4	2
Q67H/K/N	1	2
K70H/N/R/S	1	3
N74D/H/S	3	0
A105S/T	3	1
T107A/C/N	1	3

[†]Capsid genotypic and phenotypic resistance testing performed on any participants with confirmed HIV-1 RNA ≥50 c/mL and <1 log₁₀ HIV-1 RNA reduction from Day 1 at Week 4 visit, at any visit after achieving HIV-1 RNA <50 c/mL and rebound to ≥50 c/mL, and at any visit with >1 increase from nadir; HIV-1, protease, reverse-transcriptase, and integrase genotypic and phenotypic testing were performed if rebound or suboptimal virologic response were confirmed; [‡]Developed during maintenance period (Week 4 [n=5], Week 10 [n=2], and Week 26 [n=1]).


- No additional participants with LEN resistance were observed in the randomised cohort after Week 26
- All 8 participants with emergent LEN resistance remained on LEN
 - All 8 participants were at high risk of emergent LEN resistance: no fully active drugs in OBR (n=4) or inadequate adherence to OBR (n=4)
 - 3 participants resuppressed at a later visit: 1 without and 2 with OBR change

Changes in CD4 in Randomised Cohort (n=36)



- Randomised cohort: mean change in CD4, cells/μL (95% CI): 81 (44, 118) at Week 26; 83 (43, 122) at Week 52
- Non-randomised cohort: mean change in CD4, cells/μL (95% CI): 98 (59, 136) at Week 26

Changes in CD4 by Category in Randomised Cohort (n=36)



- LEN led to clinically meaningful improvement in CD4 cell count
- Proportion of participants with very low CD4 (<50 cells/μL) decreased from 22% (8/36) at baseline to 3% (1/36) at Week 52
- Proportion of participants with ≥200 CD4 cells/μL increased from 25% (9/36) at baseline to 60% (21/36) at Week 52

Acknowledgments

We extend our thanks to the study participants and their families, and the participating study investigators and staff. Canada: J Brunetta, B Trotter; Dominican Republic: E Koenig; France: J-M Molina, S Ronot-Bregigoeon, Y Yazdanpanah; Germany: H-J Stellbrink; Italy: A Antinori, A Castagna, F Castelli; Japan: T Shirasaka, Y Yokomaku; South Africa: M Rassozi; Spain: J Mallolas; Taiwan: C-C Hung; Thailand: A Avihingsanon, P Chetchotisakd, K Siripassom, W Ratanasuwam; USA: DS Berger, M Berhe, C Brinson, CM Creticos, GE Crofoot, E DeJesus, D Hagen, T Hodge, K Lichtenstein, JP McGowan, O Ogbuagu, O Osiyemi, GJ Richmond, MN Ramgopal, PJ Ruane, W Sanchez, S Segal-Maurer, J Sims, GI Sinclair, DA Wheeler, A Wiznia, K Workowski, C Zurawski. This study was funded by Gilead Sciences, Inc. Editing and production assistance were provided by BioScience Communications, New York, NY, funded by Gilead.

Adverse Events (excluding ISRs)*

≥10% Total in Any Grade, % (n)	Total LEN N=72
Diarrhoea	13 (9)
Nausea	13 (9)
COVID-19	11 (8)

*Serious adverse events (AEs) not related to study drug: malignant neoplasm and dizziness (n=1); abdominal pain, pancreatic mass, Clostridium difficile colitis, and angina pectoris (n=1); anal squamous cell carcinoma, proctalgia, impaired healing, and anal cancer (n=1); femoral neck fracture (n=1); COVID-19 (n=2); pneumonia (n=1); and septic shock, renal impairment, and shock (n=1). ISRs, injection-site reactions.

- Duration of follow up: median 376 d (interquartile range: 306, 501)
- 70 participants with ≥197 d of follow-up and 36 participants with ≥379 d of follow-up
- No serious AEs were related to study drug
- 1 participant had a serious AE of malignant neoplasm with a fatal outcome and not related to study drug

Incidence of ISRs Related to SC LEN*

ISR Types, %	After 1st SC Dose at Week 1 N=72	After 2nd SC Dose at Week 26 n=70	Median Duration, d
Swelling	26	13	12
Erythema	24	11	6
Pain	22	21	3
Nodule	22	11	180
Induration	11	10	118

*Only includes AEs related to LEN and excludes those not related to it.

- Mostly Grade 1 or 2 ISRs
- No Grade 4 ISRs, but 2 participants had Grade 3: 1 participant with swelling and erythema, which resolved in 4 and 8 d, respectively, and 1 participant with pain, which resolved in 1 d
- All nodules were Grade 1, except in 1 participant (Grade 2) who had 2 AEs of Grade 2 nodules, each after the 2nd and 3rd injections (both resolved after 3 d)
- 1 participant discontinued study drug at Week 52 due to an ISR (nodule; Grade 1)

Grade 3 or 4 Laboratory Abnormalities

Laboratory Abnormality, % (n)	Total: N=72
Any Grade 3 or 4	29 (21)
≥5% in total	
Low creatinine clearance (eGFR)*	14 (10)
Elevated creatinine [†]	13 (9)
Glycosuria	6 (4)
Nonfasting/fasting hyperglycemia	6 (3)

*Per Division of AIDS scale, Grade 3 creatinine clearance is <30 mL/min or 30–50% decrease from baseline; [†]Grade 3 creatinine is >1.8–3.5 x upper limit of normal or increase to 1.5–2.0 x baseline. eGFR, estimated glomerular filtration rate.

- None of the Grade 3 or 4 laboratory abnormalities were clinically relevant
- Low creatinine clearance/eGFR and high creatinine were transient or unconfirmed abnormalities
- Hyperglycaemia and glycosuria were transient, unconfirmed, or related to underlying diabetes

Conclusions

- In highly treatment experienced people with HIV with limited treatment options due to MDR:
 - LEN in combination with an OBR led to high rates of virologic suppression at Week 52 (83%)
 - LEN led to clinically meaningful increases in CD4 counts at Week 52
 - LEN was well tolerated, with only 1 ISR leading to discontinuation
- These data support the ongoing evaluation of LEN for treatment and prevention of HIV-1 infection
 - In highly treatment experienced people with MDR HIV
 - In treatment-naïve and -experienced people with HIV in combination with other agents
 - In people who could benefit from pre-exposure prophylaxis

References

- Link JO, et al. Nature 2020;584:614-8;
- Zila V, et al. Cell 2021;184:1032-46.e18;
- Margot N, et al. Antimicrob Agents Chemother 2021;65:e02057-20;
- VanderVeen L, et al. CROI 2021, oral 128;
- Yant SR, et al. CROI 2019, poster 480;
- Marcelin AG, et al. J Antimicrob Chemother 2020;75:1588-90;
- Gupta SK, et al. IAS 2021, oral OALB0302;
- Segal-Maurer S, et al. CROI 2021, oral 127;
- Molina JM, et al. IAS 2021, oral OALX01LB02;
- Margot N, et al. EACS 2021, oral OS1/1.
- Ogbuagu O, et al. CROI 2022, Poster 1047