

Feasibility, consumer acceptability and behavioural outcomes associated with take-home fentanyl test strips.

Rachel Sutherland¹, Maureen Steele², Edmund Silins³, Rosie Gulliver³, Phillip Read³, Amy Peacock^{1,4}, Monica Barratt^{5,1}, Robert Page^{2,1}, Raimondo Bruno^{4,1}, Craig Rodgers²

¹National Drug and Alcohol Research Centre, UNSW Sydney, Sydney, Australia; ²St Vincent's Hospital, Sydney, Australia; ³Kirketon Road Centre, South Eastern Sydney Local Health District, Sydney, Australia; ⁴School of Psychological Sciences, University of Tasmania, Hobart, Australia; ⁵Social and Global Studies Centre and Digital Ethnography Research Centre, RMIT University, Melbourne, Australia

Presenter's email: rachels@unsw.edu.au

Introduction and Aims: Fentanyl and fentanyl analogues pose an urgent public health threat. These substances are of higher potency than other opioids and are increasingly being used to adulterate heroin, contributing to tens of thousands of deaths worldwide. This project aimed to examine the feasibility, consumer acceptability, and behavioural outcomes associated with take-home fentanyl-test strips in Sydney, Australia.

Design and Methods: Seventy-eight people who had used heroin in the past six months were recruited from Kirketon Road Centre and Rankin Court, Sydney (n=40 and n=38, respectively). Participants were provided with training on how to use, and interpret the results of, fentanyl test strips. Upon completion of the training, participants were given 10 fentanyl test strips to take home for personal use and were followed up four weeks later.

Results: To-date, 67 out of 78 follow-up interviews have been completed (86%). Of those who completed the follow-up survey, 81% (54/67) reported that they had used at least one of the fentanyl test strips given to them: heroin was the most tested substance (76%), followed by methamphetamine (24%). Approximately two-fifths (42%; 22/53) reported getting at least one positive detection for fentanyl; of these, 36% (8/22) reported consuming a 'tester' prior to consumption, with smaller numbers reporting that they had gone 'slower' or used less than originally intended. Almost all participants reported they would use the strips again if they were free to access (96%) and would recommend them to peers (98%).

Discussion and Conclusions: The uptake and consumer acceptability of take-home fentanyl test strips was relatively high among our sample, providing support for its implementation and expansion across Australia.

Disclosure of Interest Statement: *AP has received untied educational grants from Seqirus and Mundipharma. RS has received untied educational grants from Seqirus. RB has received untied educational grants from Mundipharma and Indivior. Funding from these organisations has now ceased, funding was for work unrelated to this project, and the funding bodies had no role in study design, analysis and reporting. No pharmaceutical grants were received for this study.*