

# Charge equilibration times for slow highly charged ions in single layer graphene

E. Gruber<sup>\*</sup>, R. A. Wilhelm<sup>†</sup>, J. Schwestka<sup>\*</sup>, V. Smejkal<sup>\*</sup>, R. Kozubek<sup>‡</sup>, A. V. Krasheninnikov<sup>†</sup>,  
M. Schleberger<sup>‡</sup>, S. Facsko<sup>†</sup>, and F. Aumayr<sup>\* 1</sup>


<sup>\*</sup>TU Wien, Institute of Applied Physics, 1040 Vienna, Austria, EU

<sup>†</sup>Helmholtz-Zentrum Dresden-Rossendorf (HZDR), 01328 Dresden, Germany, EU

<sup>‡</sup>Universität Duisburg-Essen, 47048 Duisburg, Germany, EU


**Synopsis** We report on charge exchange and energy loss measurements for slow highly charged Xe<sup>q+</sup> ( $q \leq 35$ ) ions after transmission through a single layer of freestanding graphene. Surprisingly short charge equilibration times of only a few femtoseconds are found, which cannot be explained within currently available models.

Highly charged ions can be used as a tool to measure the short-time response of 2D materials to an extremely large, local external field. A charge state of  $q = 35$  implies a local electric field strength of  $1.8 \times 10^{11} \text{ Vm}^{-1}$  at a distance of  $5 \text{ \AA}$  from it (corresponding to laser power densities of  $\sim 10^{17} \text{ Wcm}^{-2}$ ). Previous work on scattering of HCI from solid surfaces or their transmission through freestanding carbon membranes reported unexpectedly large charge capture within 5 - 30fs [1 and refs. therein]. Recently we took the final step and presented results for the ultimately thin carbon target, a freestanding single layer of graphene [2]. Measurements of the charge state and energy of the transmitted ions surprisingly show that a large number of electrons is extracted from a small surface area (see figure 1).



**Figure 1.** The strong electric field of a highly charged ion is able to capture dozens of electrons from the single layer graphene within a few femtoseconds [2].

Charge equilibration times are derived from the mean exit charge state in dependence of the incident charge state and ion velocity. We find that Xe<sup>q+</sup> ions equilibrate within a time of only a few fs (see figure 2).



**Figure 2.** Number of captured (and stabilized) electrons after transmission of highly charged Xe ions (incident charge state  $q_{in} = 20-35$ ) through a single layer graphene sheet as a function of the inverse projectile velocity [2].

Our current understanding of HCI-solid interaction is not able to explain an almost complete de-excitation of the projectiles within the limited time span of the interaction with a 2D target. This strongly points to a new mechanism, which will be presented at the conference.

## References

- [1] R.A. Wilhelm *et al.* 2014 *Phys.Rev.Lett.* **112**, 153201
- [2] E. Gruber *et al.* 2016 *Nature Comm.* **7**, 13948

<sup>1</sup> E-mail: [aumayr@iap.tuwien.ac.at](mailto:aumayr@iap.tuwien.ac.at)