

Photodissociation of cold RbBa^+ ions produced in a hybrid cold atom-ion trap

Humberto da Silva Jr ^{*}, Maurice Raoult ^{*}, Amir Mohammadi [†], Johannes Hecker Denchlag [†],
and Olivier Dulieu ^{*1}

^{*} Laboratoire Aimé Cotton, CNRS, Univ. Paris-Sud, ENS Paris-Saclay, Univ. Paris-Saclay, 91400 Orsay, France

[†] Institut für Quantenmaterie and Center for Integrated Quantum Science and Technology, Universität Ulm, Germany

Synopsis The formation and destruction processes of cold RbBa^+ molecular ions are modelled in the context of their experimental observation in hybrid setups merging cold atom and cold ion traps.

Investigating atom-ion interactions in hybrid setups merging a cold atom trap and a cold ion trap has revealed to be one of the novel developments of research on ultracold matter. Beside elastic or inelastic binary collisions, charge exchange, or three-body collisions, the formation of cold molecular ions has been directly observed by one group [1], while they are predicted to be created in most experiments of this kind [2].

Here we present a theoretical investigation about the possible destruction of RbBa^+ molecular ions by a photodissociation mechanism, due to the many lasers involved in the experimental setup of Ref. [3]. The potential energy curves (PEC) and all the relevant permanent and transition electric dipole moments (PEDM and TEDM, respectively) of RbBa^+ are calculated in a semiempirical methodology which has been already used in our group to describe a number of diatomic systems [4]. Both atoms are represented as an one-electron system, where the valence electron is moving in the field of a relativistic effective core potential (ECP) with a core polarization potential (CPP) correction. The RbBa^+ electronic structure is thus reduced to an effective two-electron problem. The remaining two valence electrons are considered to calculate the Hartree-Fock and the excitation determinants, in an atom-centered Gaussian basis set, through the usual self-consistent field methodology. Finally, a full configuration interaction (FCI) is achieved.

With these data we computed the state-to-state absorption cross section, $\sigma_{\Lambda v j}(\nu)$, of a RbBa^+ rovibrational level (v, j) in the Λ electronic state induced by a laser with frequency ν . Following Ref. [3] we focused on photodissociation by a 1064 nm laser towards accessible electronic states Λ' , reaching continuum states with angular momentum j' and energy $\varepsilon = \hbar k$. The corresponding process is illustrated in Figure 1.




Figure 1. Computed RbBa^+ potential energy curves and transition dipole moments. The suggested transition (black arrow) corresponds to an initial weakly-bound molecular ion assumed to be created by three-body collision (see Ref. [3])

Preliminary results indicate that the lifetime of the molecular ions is strongly limited by the presence of laser light in the experiment, in agreement with the findings of Ref. [5].

References

- [1] F. H. J. Hall *et al.*, 2015 *Phys. Rev. Lett.* **107**, 243202 (2011)
- [2] H. da Silva Jr *et al.* 2015 *New J. Phys.* **17**, 045015
- [3] A. Krüchow *et al.* 2016 *Phys. Rev. A* **94** 030701 (R)
- [4] M. Aymar *et al.* 2011 *J. Chem. Phys.* **135** 064305
- [5] S. Jyothi *et al.* 2016 *Phys. Rev. Lett.* **117**, 213002

^{*} E-mail: olivier.dulieu@u-psud.fr