

Magic wavelengths of Ca⁺ ion for linearly and circularly polarized light

Jun Jiang^{*1}, Li Jiang^{*}, Xia Wang^{*}, Peter Shaw[†], Deng-Hong Zhang^{*}, Lu-You Xie^{*}
and Chen Zhong Dong^{*2}

^{*}Key laboratory of Atomic and Molecular Physics & Functional Materials of Gansu Province, College of Physics and Electronic Engineering, Northwest Normal University, Lanzhou, 730070, China

[†]School of Engineering, Charles Darwin University, Darwin NT 0909, Australia

Synopsis The dynamic dipole polarizabilities of the low-lying states of Ca⁺ ion for linearly and circularly polarized light are calculated by using relativistic configuration interaction plus core polarization (RCICP) approach. Present magic wavelengths for linearly polarized light agree with the available results excellently. Additional magic wavelengths have been found for circularly polarized light, which is very useful for magnetic-sublevel selective trapping.

The magic wavelength, at which the two levels of transitions have same AC stark shifts, was introduced in Refs. [1, 2]. It has been of great interest in ultraprecise optical lattice clocks and the state-insensitive quantum engineering [3, 4]. For circularly polarized light, the dynamic polarizability has vector and tensor component. The dynamic polarizability is different for different magnetic sublevels of atomic states.

In this work, the dynamic dipole polarizabilities of the $4s$, $4p_j$ and $3d_j$ states of Ca⁺ ion for linearly and circularly polarized light are calculated by using relativistic configuration interaction plus core polarization (RCICP) approach. The basic strategy of the model is to partition the atom into valence and core electrons. The orbitals of the core are written as linear combinations of S-spinors which can be treated as relativistic generalizations of Slater-type orbitals. The orbitals of valence are written as linear combinations of L-spinors and S-spinors. L-spinors can be treated as relativistic generalizations of the Laguerre- type orbitals.

Fig.1 shows the dynamic polarizabilities of $4s$ and $4p_{1/2}$ states for the linearly ($A=0$) and left handed ($A=-1$) polarized light. In Fig.1 (a), three magic wavelengths are found, which are in good agreement with available results [5, 6]. Fig.1 (b) gives the dynamic polarizabilities for $A = -1$ for $4s_{m=-1/2}$ and $4p_{1/2m=-1/2}$ states. There are only two magic wavelengths are found for $4s - 4p_{1/2}$ transition. There is no magic wavelength between $4p_{1/2} - 5s$ and $4p_{1/2} - 4d_{3/2}$ transition energy. The magic wavelength 395.5410 nm very close to the magic wavelength 395.1788 nm for $4s - 4p_{1/2}$ transition for $A=0$. Another magic wavelength 778.3726 nm occurs between $4p_{1/2} - 4s$ and $4p_{1/2} - 3d_{3/2}$ transition energy. It has 87 nm difference with the magic wavelength 691.2444 nm for linearly polarized light.




Fig. 1. Dynamic polarizabilities (in au) for the $4s - 4p_{1/2}$ of Ca⁺ for the linearly ($A=0$) and left handed ($A=-1$) polarized light. Magic wavelengths are identified by arrows.

This work was supported by National Natural Science Foundation of China (Grant No. 11564036, U1332206, 11464042, U1330117, U1331122, U153014), the Young Teachers Scientific Research Ability Promotion Plan of Northwest Normal University (NWNLU-LKQN-15-3).

References

- [1] J. Ye *et al.* 1999 *Phys Rev. Lett.* 83,4987
- [2] H. Katori *et al.* 1999 *Phys Rev. Lett.* 82,1116
- [3] G. Wilpers *et al.* 2007 *Metrologia* 44,146
- [4] N. Lundblad *et al.* 2010 *Phys Rev.A* 81,031611
- [5] J. Kaur *et al.* 2015 *Phys Rev.A* 92,031402
- [6] Y. B. Tang *et al.* 2013 *Phys Rev.A* 87,042517

¹E-mail: phyjiang@yeah.net

²E-mail: dongcz@nwnu.edu.cn