

Determination of the ${}^4P^o/{}^2P$ ratio in single electron capture of C^{4+} ($1s^2, 1s2s\ ^3S$) mixed state ion beams in 6-18 MeV collisions with H_2 , He, Ne and Ar targets

I. Madesis^{**}, A. Laoutaris^{**}, E. P. Benis[‡], T. Kirchner[♦], T. J. M. Zouros^{1**}

^{*}Department of Physics, University of Crete, P.O. Box 2208, GR 71003 Heraklion, Greece

[†]Tandem Accelerator Laboratory, INPP, NCSR Demokritos, GR 15310 Ag. Paraskevi, Greece

[‡]Department of Physics, University of Ioannina, GR 45110 Ioannina, Greece

[♦]Department of Physics and Astronomy, York University, Toronto, Ontario, M3J 1P3 Canada

Synopsis: Zero-degree Auger projectile electron spectroscopy of pre-excited He-like ions colliding with various gas targets is used with a new experimental technique which relies on the combination of gas and foil strippers to vary the metastable fraction sufficiently to allow for the determination of the ratio ${}^4P^o/{}^2P$ of recent interest. Results to date for C^{4+} ions will be presented.

A zero-degree Auger projectile spectroscopy apparatus composed of a single stage hemispherical deflector analyzer with four-element injection lens and a position sensitive detector combined with a doubly differentially pumped gas target has been newly set up for high resolution studies of electrons emitted from ions colliding with gas targets. Using this setup we have started a systematic isoelectronic investigation of projectile K-Auger electrons emitted from pre-excited He-like ions in collisions with gas targets [1]. The goal is to study single electron capture to the $1s2s\ ^3S$ long-lived component of the typically mixed state ($1s^2, 1s2s\ ^3S$) He-like ion beam. The observed KLL Auger lines for 9 MeV C^{4+} collisions, shown in Fig. 1, can be produced either by direct electron capture to the $1s2s\ ^3S$ component or by transfer-excitation from the $1s^2$ ground state [2-4]. The ratio R_m of ${}^4P^o/{}^2P$ line intensities due to capture to the $1s2s\ ^3S$ has been found to be greatly enhanced from the spin statistics [2-5] predicted value of 2, invoking contentious explanations. To isolate the $1s2s\ ^3S$ capture contributions, a new technique is employed [5] requiring the measurement of two Auger KLL spectra using C^{4+} ions with sufficiently different $1s2s\ ^3S$ metastable fraction. This fraction can be varied by the judicious use of terminal and/or post (terminal) foil or gas stripping [5].

The measured R_m values depend critically on the solid angle correction of the 4P_j yields. These states are long-lived, Auger decaying all along the projectile path, after excitation in the target gas cell. This correction has been determined and applied to our measurements using a new Monte Carlo electron trajectory simulation approach [6] with the use of the SIMION charged particle optics software [7].




Figure 1. $C^{3+}(1s2l2l')$ Auger KLL spectra from 9 MeV C^{4+} collisions with various gas targets for different $1s2s\ ^3S$ metastable beam fractions depending on gas terminal stripping (GTS) or gas post-stripping (GPS). The 4P line intensity is proportional to the $1s2s\ ^3S$ beam content.

References

- [1] I. Madesis et al. 2015 *J. Phys: Conf. Ser.* **583** 012014
- [2] D. Strohschein et al. 2008 *Phys. Rev. A* **77** 022706
- [3] T. J. M. Zouros, B. Sulik, L. Gulyas, and K. Tökési 2008 *Phys. Rev. A* **77** 050701
- [4] D. Rohrbein, T. Kirchner, and S. Fritzsche 2010 *Phys. Rev. A* **81** 042701
- [5] E. P. Benis and T. J. M. Zouros 2016 *J. Phys. B* **49** 0235202
- [6] S. Doukas et al. 2015 *Rev. Sci. Instrum.* **86** 043111
- [7] SIMION 8.1, SIS Inc. Ringoes, NJ.

¹E-mail: tzouros@physics.uoc.gr