

Effects of Regional Meteorological and Air Conditions on Community-Acquired Pneumonia -

Examining the Interaction of Individual, Meteorological, and Air Characteristics

Thomas Brenner, Ann-Christine Link, Christoph Reudenbach, Hendrik Pott, Jan Rupp, Martin Witzenrath, Gernot Rohde, Mathias Pletz, Wilhelm Bertrams, Bernd Schmeck, CAPNETZ study group

Abstract

Climate change is not only associated with both increases in frequency and intensity of extreme weather events but also elevated climate variability. As a result, weather conditions favouring pneumonia occurrence can increase, leading to adverse effects on peoples' health (Sohn et al., 2019). Community-acquired pneumonia, in the following called pneumonia, is one of the most common causes of death worldwide (Aliberti et al., 2021). At the same time, clear linkages between this disease and both meteorological and air conditions are present (Wang et al., 2021).

It is well studied that especially extreme weather and air conditions, the latter including air quality and wind conditions, have an impact on the number of people hospitalized for pneumonia (Y. Liu et al., 2014; Onozuka et al., 2009). However, many studies analyse the effects of meteorological and air conditions separately with mortality data as an endpoint while predominantly covering the Asian continent and specifically larger cities (Basu & Samet, 2002; Chung et al., 2009; Ge et al., 2013). Therefore, data on continuous geographical regions and combinations of rural and metropolitan areas are required. Additionally, little is known about how personal characteristics (age, sex) and health background (smoking history, chronic lung diseases, heart insufficiency, overweight) affect the sensitivity of pneumonia cases regarding meteorological and air conditions.

We close this research gap by analysing a prospective multicenter cohort that was treated in an in- or outpatient setting for pneumonia in 22 German hospitals and outpatient clinics between 2003-2017. The dataset contains personal and health information for more than 10,000 patients. We match this data with daily regional meteorological and air condition data while not only considering the conditions on the day of hospitalization but also up to four days before. Logistic regressions are used to examine the impact of meteorological and air conditions on pneumonia cases considering a short-term and long-term perspective as well as the modification effects of various personal characteristics on these relationships. Therefore, two different statistical approaches are used: first, the total comparison to determine under what weather and air conditions pneumonia occurs more often and second, contemporary comparison to examine why pneumonia occurs exactly on a specific and not any other day. This research is valuable since it aims to not only determine but also predict when certain groups of people are at increased risk of pneumonia. This can support health care providers during periods of weather conditions which promote pneumonia in better preparing resources such as staff and treatments but also in guiding prophylaxis such as limiting outdoor activities.

Keywords: pneumonia, climate change, meteorology, air conditions, Germany.

Contents

Abstract	1
Contents.....	2
1. Introduction	4
2.1 Methodology	6
2.1 Medical data description	6
2.2 Meteorological and air quality data description	7
2.3 Methodological approach	9
3. Results	11
3.1 Analysis of all cases	11
3.2 Influence of personal characteristics	13
References	19

Table of Figures

Figure 1: Hospitals and Outpatient Clinics with 30-minute and 60-minute catchments.	7
Figure A2: Pneumonia Deaths per 100,000 inhabitants (age-standardized): Germany.	23
Figure A3: Number of people hospitalized for pneumonia: Germany.	23

Table 1: Meteorological and Air Conditions included in the study.	8
Table 2: Regression results for the two created databases (p-values in brackets).	12
Table 3: Regression results for contemporary comparison including interaction terms with personal characteristics (p-values in brackets).	14
Table 4: Regression results for total comparison including interaction terms with personal characteristics (p-values in brackets).	15

Acronyms

AIC: Akaike Information Criterion

BMI: Body Mass Index

CO: Carbon monoxide

COPD: chronic obstructive pulmonary disease

CAMS: Copernicus Atmosphere Monitoring Service

DTR: Diurnal Temperature Range

DWD: German Weather Service (Deutscher Wetterdienst)

IQTIG: Institut für Qualitätssicherung und Transparenz im Gesundheitswesen

NO₂: Nitrogen dioxide

RH: Relative Humidity

RKI: Robert Koch Institut

SO₂: Sulfur dioxide

O₃: Ozone

PM: Particulate matter

PT: Pneumonia Temperature

SRTM: Shuttle Radar Topography Mission

WHO: World Health Organization

WMO: World Meteorological Organization

1. Introduction

Respiratory diseases, including pneumonia, have a high societal and economic importance not just worldwide but also in Germany (Plass et al., 2014; World Health Organization, 2007). In Germany in 2008, the treatment of patients with respiratory diseases led to treatments costs of 14.2 billion euros (Gillissen & Welte, 2014). The yearly number of community-acquired pneumonia cases in Germany is estimated to lie between 400,000 and 600,000 (RKI, 2017). About 30 to 50 % of these pneumonia cases end up being treated in hospitals (Höffken et al., 2009; Schnoor et al., 2007). The age-related mortality of pneumonia in Germany is 13% (Institut für Qualitätssicherung und Transparenz im Gesundheitswesen [IQTiG], 2016). At the same time, this number is expected to be even larger due to underestimations associated with the monocausal nature of mortality statistics.

Apart from the societal and economic effects of pneumonia, climate change and its impact on health in general but pneumonia specifically has gained broad scientific attention over the last decades. It has been well studied that especially extreme weather and air conditions, the latter including air quality and wind factors, affect the number of people hospitalized for pneumonia (Y. Liu et al., 2014; Onozuka et al., 2009; Wang et al., 2021). Climate change is associated with not only more frequent and intense weather extremes but also a higher variability in weather conditions which can negatively affect peoples' health (IPCC, 2022). In the next sections, we will present different studies which evaluated the possible relationship between meteorological and air conditions on pneumonia and other well-known respiratory diseases.

Temperature was determined to be one of the factors that shows the greatest contribution to pneumonia mortality and morbidity (Braga et al., 2002; Kim et al., 2016; L. Liu et al., 2011). However, the pattern of the relationship between temperature and pneumonia (j-shaped, u-shaped or v-shaped) strongly depends on the study location and associated climate (Braga et al., 2002; Falagas et al., 2008; Nascimento-Carvalho et al., 2010) as well as the selected temperature indicator (average, minimum or maximum temperature, pneumonia temperature, diurnal temperature range etc.). Nonetheless, studies predominantly detected a u-shape pattern, indicating that not only minimum but also maximum temperatures are causing higher numbers of pneumonia cases, especially in the elderly subgroup (Green et al., 2010; Y. Liu et al., 2014). The most widely used temperature factors include the diurnal temperature range (DTR) defined as the difference between the minimum temperature and maximum temperature on a given day and pneumonia temperature (PT), defined as the current day's average temperature minus the previous day's average temperature. The majority of studies determined that increases in diurnal temperature correlate with increasing pneumonia cases (Lim et al., 2012; Xu et al., 2013). At the same time, many studies also found that pneumonia is more likely during colder periods, for example, during winter (Eccles & Wilkinson, 2015; Mäkinen et al., 2009).

Relationships of both precipitation and relative humidity (RH) with pneumonia have predominantly been reported in countries with abundant rainfall and warmer average temperatures (Chan et al., 2002; Mäkinen et al., 2009; Nascimento-Carvalho et al., 2010). Furthermore, Onozuka et al. (2009) found that increasing relative humidity correlates with elevated numbers of pneumonia cases since the survival of the pneumonia virus in aerosols was found to be best at extremes of relative humidity.

Moreover, some studies account for personal characteristics predominantly sex and age when assessing the correlation of meteorological and air conditions with pneumonia cases (Lam et al., 2018; Sohn et al., 2019). Some also account for one or two diseases (health background) of pneumonia patients in their study (Basu & Samet, 2002). For example the research by Basu and Samet (2002) indicates that people with pre-existing cardiovascular and respiratory diseases have an elevated risk of death from pneumonia associated with especially ambient heat exposure and that risk is also higher for specific subgroups such as the elderly, infants, and persons of low socioeconomic status.

Studies that assess the relationship of sunshine and wind speed with pneumonia cases are rare. For example, Lam et al. (2018) detects increased pneumonia hospitalization in the elderly (>75 years) during high wind conditions. Paynter et al. (2013) detects a strong association of sunshine with pneumonia; however their study was conducted in the tropics which are associated with a completely different climate and temperature regime than Germany which has a temperate climate. Moreover, Murdoch and Jennings (2009) analysed the association of respiratory virus activity and environmental factors with the incidence of invasive pneumococcal disease (instead of pneumonia). They found significant correlations between pneumococcal cases and decreasing sunshine hours and wind speed, and with increasing air pollution levels and humidity.

Regarding air quality, the relations among air pollutant levels, temperature, and predominantly pneumonia mortality have been investigated in several studies; however, leading to conflicting results. For example, the literature review of the short-term correlation between ambient air pollutants (sulfur dioxide (SO₂), ozone (O₃), nitrogen dioxide (NO₂), carbon monoxide (CO), particulate matter (PM) with diameter 10 (PM₁₀) and 2.5 mm (PM_{2.5})) and pneumonia hospitalization of children by Nhung et al. (2017) concludes that the ambient air pollution, especially O₃, NO₂, CO, PM₁₀, PM₂₅ and SO₂, organized in decreasing impact, was related to both pneumonia cases and severity. Also, a long-term prospective, observational study in a Chinese hospital showed that meteorological and air pollution conditions contributed more to severe pneumonia in children than respiratory viruses did (Wang et al., 2021). Moreover, Chiu et al. (2009) also examined that high temperatures (>23°C) and elevated levels of SO₂ are correlated with increasing numbers of pneumonia cases. Nevertheless, others such as Katsouyanni et al. (1997) see air pollutants rather as “confounders and/or effect modifiers” (i.e., a synergistic effect) of the temperature-mortality association while no effect is not found by others such as Samet et al. (1998).

As presented above, many studies attempted to examine the relationship between meteorological and health conditions with pneumonia while placing a focus on specific meteorological or air conditions in relatively broad areas, predominantly Asia, with mainly mortality data as an endpoint (Basu & Samet, 2002; Chung et al., 2009; Ge et al., 2013). However, health issues should not solely be based on outcomes such as mortality but also morbidity. Furthermore, only scarcely did studies account for how the sensitivity of pneumonia cases to meteorological and air conditions differs when considering personal characteristics and especially health background.

We address the previously identified shortcomings by not only including the personal characteristics of 10,000 patients treated in 22 German hospitals or outpatient clinics for pneumonia but also health background information such as smoking history, cardiac insufficiency, chronic pulmonary disease, diabetes, and overweight in our analysis. We match this data with high daily spatial and temporal resolution, regional meteorological and air conditions data, only considering data in a one-hour catchment surrounding each hospital. The included regional meteorological and air conditions are displayed in Table 1. Logistic regressions are then used to examine the effects of meteorological and air conditions on pneumonia cases. Furthermore, we want to disentangle how the sensitivity of pneumonia cases to meteorological and air conditions changes when considering the patients' personal and health backgrounds.

This study adds new insights to the discussion by taking two statistical approaches to the regression (which will be explained in more detail in section 2.3): 1) the total comparison approach aims to determine under which meteorological and air conditions pneumonia occurs more frequently and 2) the contemporary comparison approach aims to determine which factors potentially cause pneumonia to occur on specific days and not others. So far, the studies in this field tend to follow a total comparison approach, aiming to find general meteorological and air conditions which are more likely to appear in accordance with pneumonia. We aim to quantify the relationship between both meteorological and air conditions with pneumonia cases to not only determine but predict when certain groups of people are at an increased risk of pneumonia. This would both support health care providers in better preparing resources such as staff and treatments and guiding prophylaxis such as limiting outdoor activities during periods of meteorological conditions which promote pneumonia.

We find that several meteorological and air conditions influence the risk of pneumonia not on their own but specifically in combination with other factors: e.g., low temperature in combination with low air pressure or high air pressure together with high air pollution lead to more pneumonia cases. Furthermore, we find several differences in the sensitivity of various population groups to the different meteorological and air conditions.

2.1 Methodology

2.1 Medical data description

The medical data contains information on 10,808 people admitted for pneumonia from 22 German hospital or doctor's practices between February 2003 and December 2017 including the exact day of hospitalisation (Figure 1). The following diseases are accounted for in the health background of each pneumonia patient: history of smoking, body mass index (BMI), tumour, diabetes, cardiac insufficiency, and chronic pulmonary disease.

Hospitals and Outpatient Clinics with 30-minute and 60-min catchments




Figure 1: Hospitals and Outpatient Clinics with 30-minute and 60-minute catchments.

2.2 Meteorological and air quality data description

The climate data used in this study comes from the operational official station monitoring networks of the German Weather Service (DWD). This data covers the period from 2000 until 2021 and includes all daily available meteorological station data (Behrendt et al., 2011; DWD Climate Data Center, 2021). Within the study period, a larger number of stations was converted to automatic data acquisition in 2009, which led to data outages in some cases. Quality controls include temporal and content consistency checks to correct for random and gross measurement errors, following the World Meteorological Organization (WMO) and DWD standards (Kaspar et al., 2013). The daily mean values of air pressure, relative humidity, temperature, minimum maximum temperature, and the number of hours of sunshine come from the previously described data source.

The Shuttle Radar Topography Mission (SRTM) terrain model, version 4, is used to account for diurnal climatological effects due to terrain. Data with an average resolution of 60x60 meters in Germany is resampled bilinearly to 500*500-meter cells (Jarvis et al., 2008; Reuter et al., 2007).

To obtain regionalized climate values for each municipality in Germany, it is necessary to spatially interpolate the daily climate data since it is only available as point measurements. Numerous approaches exist to perform such interpolations, some of which differ considerably. Besides comparatively simple approaches like an inverse distance weighted method or spline-based methods, the kriging methods have proven to be very useful technique for interpolations. In situations with a changing and small number of point data (climate stations), ordinary kriging with external drift (KED) is the most efficient standard interpolation method and is the most

commonly used geostatistical approach for estimating climate variables (Benavides et al., 2007). Since the above-mentioned meteorological variables are known to be dependent on topography, KED is used considering terrain elevations as secondary data to estimate the climate variables of interest (Bianchi et al., 2016).

In the kriging-based approaches, the study area of Germany is divided into a finite number of cells (2.226.005) with a resolution of 500 m to determine climate parameter fields. A two-step procedure is then applied to compute daily meteorological maps using R software. In the first step, an autoadaptive optimization of the variogram data is determined from the co-variance models (Gräler et al., 2016; Pebesma, 2004; R Core Team, 2021). In the second step, the optimized covariance model is utilized for each daily meteorological variable in the universal kriging procedure. Kriging differs from simpler interpolation methods in that it is based on the spatial arrangement of the empirical observations rather than on the assumed model of spatial distribution. With at least moderate spatial autocorrelation, it is a useful method for preserving spatial variability. The universal kriging used in this study has a modified assumption of stationarity in that the mean of the values at different locations can be deterministically different while only the variance remains constant across the field (Bivand et al., 2008). This second-order stationarity is often an important assumption for environmental exposure such as climate values or pollution levels. To represent the external drift, the SRTM terrain heights are used and integrated into the Kriging procedure.

The data on total precipitation and wind speed comes from a different data set, namely from the COSMO-REA6 reanalysis, with a grid resolution of 6m x 6km for 2000 until August 2019 covering Europe (Bollmeyer et al., 2015). The extreme precipitation data is based on E-OBS observational data at the daily temporal scale and a spatial resolution of 0.02° x 0.02° (Copernicus Climate Change Service, 2020). The air pollutant data comes from Copernicus Atmosphere Monitoring Service (CAMS) global reanalysis (EAC4) with a spatial resolution of 0.75° x 0.75° (Inness et al., 2019).

As a final step, current municipality geometries were homogenized to 2012 based on official EuroStat data (Europäische Kommission - Eurostat/GISCO, 2022) to then calculate descriptive statistics of climate variables for each of the municipalities. Proportional grid cells with appropriate precision weighting were also considered.

Table 1: Meteorological and Air Conditions included in the study.

Factor	Unit	Description
Sunshine duration	hours	Daily hours of sunshine
Air pressure	Pa	Average daily air pressure
Relative Humidity	K	Daily average relative humidity
Average, minimum, and maximum temperature	°C	Daily average, minimum, and maximum temperature

Diurnal temperature difference	°C	Daily difference between the minimum temperature and maximum temperature on a given day
Pneumonia temperature	°C	Current day's average temperature minus the previous day's average temperature
Total precipitation	kg/m ²	Daily total precipitation
Precipitation at fixed percentiles (95 th , 99 th percentile)	kg/m ²	Total precipitation when daily precipitation amounts exceed the 90th, 95th, or 99th percentiles in wet days (daily precipitation \geq 1 mm) computed over the 30-year period (1989-2018).
Maximum wind speed	m/s	Daily 10 m maximum wind speed
Maximum total column* CO	kg kg ⁻¹	Daily maximum total column of CO
Total column* NO ₂	kg kg ⁻¹	Daily maximum total column of NO ₂
Total column* O ₃	kg/m ²	Daily maximum total column of O ₃
Total column* SO ₂	kg kg ⁻¹	Daily maximum total column of SO ₂
PM 10 concentrations	kg/m ³	Daily maximum of particulate matter < 10 μ m (aerosol)
PM 2.5 concentrations	kg/m ³	Daily maximum of particulate matter < 2.5 μ m (aerosol)
Black carbon aerosol optical depth**	dimensionless	Daily maximum aerosol optical depth at 550 nm for black carbon
Dust aerosol optical depth**	dimensionless	Daily maximum aerosol optical depth at 550 nm for dust aerosols
Total aerosol optical depth**	dimensionless	Daily maximum aerosol optical depth at 550 nm for total aerosols

* Total column refers to the total amount of the selected variable in a column of air extending from the surface of the Earth to the top of the atmosphere.

**The aerosol optical depth measures the amount of light lost due to the presence of aerosols on a vertical path through the atmosphere.

2.3 Methodological approach

To assess the impact of meteorological and air conditions on pneumonia morbidity, we generate for each person in the medical database three alternative dataset entries that represent all personal characteristics except the hospitalisation date. These can be interpreted as alternative days at which the person could have been admitted but actually has not been admitted. As a

result, we obtain 43.232 cases, 10.808 real cases and 32.424 non-real cases. Next, we match the meteorological and air condition data to each of these cases. Logistic regressions are used to then examine the determinants of the occurrence of the real cases in comparison to the non-real cases.

In this approach the calculation of the non-real cases is crucial. Here, we applied two different procedures:

1) Contemporary comparison: In this procedure, a day is randomly drawn from the period spanning from ten days before the real case to ten days after the real case. Hence, all non-real cases fall into the same part of the year, excluding seasonal effects from the analysis. The goal pursued with this procedure is to determine the factors which potentially caused pneumonia to occur exactly on this day and not a few days earlier or later. The question asked in this analysis is: What meteorological and air conditions trigger pneumonia?

2) Total comparison: In this procedure, the days of admission for the non-real cases are drawn randomly from the whole period of observation from the respective hospital (the periods of observation are different for the considered hospitals or outpatient clinics). In this case, the results show which general meteorological and air conditions are more likely to appear together with pneumonia cases. By following this approach, the seasonal characteristics linked to pneumonia are also identified. The question asked in this analysis is: Under what meteorological and air conditions does pneumonia occur more frequently?

The adequateness of the logistic regression is tested with the Hosmer-Lemeshow-Test, which identifies a slightly significant deviation from the assumed logistic distribution ($p\text{-value}=0.0438$). The meteorological and air conditions are assigned to each real and non-real case based on the location of the hospital and the day of hospitalisation. In general, the hospitalised people do not live at the location of the hospital but probably in the surrounding area. Therefore, we calculate the meteorological and air conditions in all municipalities that are less than 60 minutes driving time away from the hospital to approximate the relevant conditions. A weighted average is built for each hospital. Each municipality is weighted according to its population (larger cities have a higher probability that the patient originates from this city) and according to its distance to the hospital, a distance decay function is used as described in Brenner and Pudenko (2019).

We include 20 meteorological and air condition variables in the analysis (Table 1). However, several conditions are highly correlated with each other and describe similar aspects. This study includes three variables for temperature (average, minimum, and maximum), three variables for precipitation (total, $>95\%$, and $>99\%$) and eight variables associated with air pollution (CO , NO_2 , O_3 , SO_2 , PM_{10} , $\text{PM}_{2.5}$, black carbon aerosol, and total aerosol). Including them together leads to multicollinearity problems. Hence, for each aspect only one variable is included, the one leading to the highest fit (lowest AIC).

Besides including the variables (and their variants) directly as independent variables, we also test the following additional versions:

- We study whether the change of variables from one day to the next has an impact (in the case average temperature this is called the Pneumonia temperature).

- For each independent variable, a quadric term is also considered. The aim here is to examine non-linear dependencies with a specific focus on u-shaped and inverted u-shaped effects.
- Furthermore, interaction terms are included in the analyses. Each pair of variables is considered to check whether the co-occurrence of conditions is specifically triggering pneumonia.

Four variants of time delays are examined: using the values on the day of hospitalisation, one, two and three days before hospitalisation. The variant is chosen according to the AIC (Akaike Information Criterion). The aforementioned approach results in 240 potential independent variables. A stepwise in- and exclusion of the variables is used, adding one variable after the other and eliminating insignificant (significance level: .05) variables. During the approach the models are checked for multicollinearity and in case of potential multicollinearity (variance inflation factor values above 10) those variables are excluded that lead to the lowest loss in model fitting (lowest AIC values).

3. Results

In the following subsections the results of the regression analysis of all cases (section 3.1) and the influence of personal characteristics on the sensitivity of pneumonia (section 3.2) are presented and discussed.

3.1 Analysis of all cases

The first interesting result concerns the time delay for the meteorological and air conditions: In the case of the total comparison approach where non-real cases are drawn randomly from the whole period, using the conditions from two days before hospitalisation, leads to the best fit. In the case of the contemporary comparison where non-real cases are drawn from nearby days within 10 days before and after, the best fit results from using the conditions from the day of hospitalisation. Above we argue that the contemporary comparison focuses on triggering conditions, which seem to be present at the day of hospitalisation. The total comparison rather focuses on more long-lasting meteorological and air conditions, so that more days matter, reflected by a time distance of two days.

A further difference between the analysis types is the finding that three of the five significant determinants are changes in the variables in the contemporary comparison, while this is the case only for two out of 16 significant variables in the total comparison (Table 3). Thus, strong changes in the meteorological and air conditions trigger pneumonia cases: a decrease in sunshine hours, an increase in the diurnal temperature (difference between minimum and maximum temperature), and an increase in air pollution (NO₂ is most significant but the other air pollution variables lead to similar results). The fourth significant trigger (contemporary comparison) is an interaction term that also represents meteorological changes: Long sunshine duration and high precipitation (>95%-percentile) occurring together in a day also lead to more pneumonia cases. The fifth variable found significant in the contemporary analysis is

interaction term between high precipitation (>95%) and dust aerosols, which is associated with the fact that intensive precipitation reduces the effects of air pollution (Kwak et al., 2017).

Table 2: Regression results for the two created databases (*p*-values in brackets).

Variable	Contemporary comparison	Total comparison
Sunshine duration		.0339*** (.0000)
Sunshine - Change	-.00828** (.0074)	
Interaction: Sunshine * Prec. (>95%)	.0314* (.0128)	
Air pressure		-.00713*** (.0000)
Air pressure - Squared		.000165*** (.0001)
Relative humidity		.00426* (.0204)
Relative humidity - Squared		-.000721*** (.0000)
Interaction: Humidity * Diurnal temp.		-.000280*** (.0000)
Diurnal temp. - Change	$6.342 \cdot 10^{-13}$ * (.0342)	
Minimum temp. - Change		.0184*** (.0002)
Minimum temp. - Squared		-.000889*** (.0000)
Interaction: Min. temp. * Air pressure		-.0000338*** (.0000)
Interaction: Min. temp. * SO ₂		.00281*** (.0000)
Interaction: Air pressure * SO ₂		.000110*** (.0000)
High precipitation (>99%) - Squared		-.6104** (.0012)
Interaction: Prec. (95%) * Dust aerosols	-.00000198** (.0061)	
Dust aerosols - Squared		$9.304 \cdot 10^{-13}$ * (.0216)
Interaction: Dust aerosols * SO ₂		-.000000164** (.0001)
Maximum SO ₂ - Change		-.00521*** (.0000)
Maximum NO ₂ - Change	.00368** (.0074)	
Maximum SO ₂ - Squared		-.00279*** (.0000)

The total comparison provides a long list of significant variables. While looking at these factors we must keep in mind that they characterise more long-lasting meteorological and air conditions as well as seasons in which more pneumonia cases occur. We find that pneumonia is more frequent in periods with a lot of sunshine, very low air pressure (due to quadratic term), and very high relative humidity. The latter effect is increased in case of small temperature difference within a day (diurnal temperature). Studies have shown that fluctuations in humidity can lead

to changes in the respiratory epithelium which elevates the susceptibility to infectious diseases such as pneumonia (Deal et al., 1980).

Interestingly we do not find a direct linear effect of temperature. The negative quadratic effect implies that pneumonia is more likely for average minimum temperatures. Furthermore, an increase in the minimum temperature comes together with more pneumonia cases. In contrast to this, a regression with only including (minimum) temperature leads to the clear result that pneumonia is more likely in colder times. A proposed mechanism to explain why cold temperatures cause respiratory symptoms is that long-term exposure to cold air temperatures induces inflammatory reactions in the lower respiratory tract (Larsson et al., 1993). Nonetheless, even if short-term exposure to cold air may elevate the number of inflammatory cells, there is a lack of evidence to conclude that solely an increase in number is sufficient to trigger pathological reactions (Larsson et al., 1998). At the same time, one also needs to consider that during periods of cold temperature, people spend more time indoors which increases the opportunities for spread of infections between people.

However, the complete regression (Table 3) shows that temperature rather influences pneumonia together with other factors. First, it increases the effect of low air pressure. If low air pressure coincides with low minimum temperatures pneumonia is especially likely. Second, high temperature, as well as high air pressures, lead together with high air pollution (represented by SO_2) to more pneumonia cases.

No clear results are found for precipitation: the quadratic dependence is difficult to interpret because the variable takes average values only if some of the surrounding area of a hospital has very high rainfall (above 99%-percentile) but not all of the surrounding area. This would make pneumonia more likely and might indicate that punctual heavy rainfall is a meteorological condition which promotes pneumonia.

Surprisingly, we find rather negative effects of air pollution on pneumonia likelihood. Especially if dust aerosols and air pollution (represented by SO_2) are both low, pneumonia cases are more frequent. However, this must be seen in the context of the other findings: Air pollution together with high temperature or with high air pressures leads to more pneumonia cases. This outweighs the effects found for the pure air condition variables, which is confirmed by a significant correlation between air pollution and pneumonia in our data. It is given that in many locations, patterns of air pollution are driven by the weather, and at the same time air pollutant concentrations may be associated with temperature. Nevertheless, most studies show that air pollution does not influence pneumonia cases on its own but becomes relevant in connection with other conditions, in our study in connection with high air pressure or high temperatures. This nonetheless shows that additional studies are needed to further examine the effect of the relationship between air pollutants and temperature on pneumonia mortality but more specifically on morbidity.

3.2 Influence of personal characteristics

To study whether the impact of meteorological and air conditions on pneumonia depends on personal characteristics, we add to the previously identified best models (with the determined lags and variable choices) addition interaction terms between all independent variables and

various personal characteristics. Again, a stepwise approach is used where insignificant interaction terms are excluded. For the contemporary comparison, the results are listed in Table 3.

While the impact of sunshine duration change (and interaction with precipitation) and air pollution change is not influenced by personal characteristics (regression results remain relatively stable), other effects are strongly influenced. The results show that effects of increases in diurnal temperature is only given for people with a high BMI (BMI>25). In addition, people with a high BMI are hospitalised for pneumonia more often if the amount of dust aerosols in the air is high or low.

Dust aerosols are found to have an impact mainly on people who smoke. This group of people is especially affected by the combination of high humidity and high levels of dust. This effect is counterbalanced by precipitation.

People with a tumour are more likely affected by pneumonia if relative humidity decreases from one day to the other. The same holds for people with diabetes who also benefit more from the counterbalancing effect of precipitation on air pollution. An interesting fact is that change in sunshine hours disappears by including the interaction term with diabetes. This implies that decreases in sunshine from one day to the next triggers pneumonia mainly for diabetes patients and not (significantly) for other people. We also tested various other personal characteristics (such as chronic pulmonary diseases and cardiac insufficiency) without significant results for the contemporary comparison.

Table 3: Regression results for contemporary comparison including interaction terms with personal characteristics (p-values in brackets).

Variable	Personal characteristic			
	Smoking	BMI>25	Tumour	Diabetes
Sunshine - Change	-.00758* (.0386)	-.00747* (.0416)	-.00952** (.0095)	Not significant
Interaction: Sun * Prec. (>95%)	.0307* (.0160)	.0306* (.0152)	.0308* (.0143)	.0319* (.0126)
Diurnal temp. - Change	not significant	not significant	not significant	not significant
Interaction: Prec. (95%) * Dust	not significant	-.00000188** (.0068)	-.00000153* (.0275)	-.00000143* (.0378)
NO ₂ - Change	.00363** (.0083)	.00375** (.0065)	.00396** (.0040)	.00402** (.0036)
<i>Personal characteristics interaction terms:</i>				
Interaction: Humidity * Dust	1.11*10 ⁻⁸ * (.0146)			
Interaction: Prec. * Dust	-.00000405* (.0240)			
Diurnal temp. - Change		.0157* (.0470)		

Dust aerosols - Squared		9.172*10 ⁻¹³ * (.0359)		
Humidity - Change			-.0293*** (.0000)	-.0120*** (.0017)
Sunshine - Change				-.0241* (.0247)
Interaction: Prec. (95%) * NO ₂				-.0120* (.0305)

Furthermore, we determined that personal characteristics matter more in the total comparison (Table 4) compared to the contemporary comparison. While we did not find any dependence on wind speed in the analyses above it seems to matter for people who smoke since they are more likely to get admitted for pneumonia if wind speed increases from one day to the next. Low wind speeds are also connected to a higher likelihood of pneumonia infections for tumour patients. Furthermore, tumour patients show an additional connection of pneumonia with higher minimum temperatures, increases in sunshine hours, and high air pollution together with high diurnal temperatures.

Table 4: Regression results for total comparison including interaction terms with personal characteristics (p-values in brackets).

	Personal characteristic						
Variable	Smoking	Tumour	BMI>25	Chronic pulmonary disease	Age>64	Sex (female)	Cardiac insufficiency
Variables as above (Table 2)	Results similar to above (Table 2)						
Minimum temp. - Squared					Not significant		
<i>Personal characteristics interaction terms</i>							
Wind speed - Change	5.71*10 ⁻¹¹ * (.0284)						
Humidity							-.00221** (.0045)
SO ₂			-.0177** (.0026)	.0242*** (.0010)			
Diurnal temp.			.00825* (.0115)				.0180** (.0031)
Air pressure - Change			.00803* (.0127)				

Interaction: Sunshine * Prec. (>99%)			-.188*** (.0008)			-.0754* (.0434)	
Interaction: Prec. (>99%) * Diurnal temp.			.109*** (.0002)				
Minimum temp. - Squared					-.000866* (.0209)		
Minimum temp.		.0159* (.0126)			.0133** (.0026)		
Wind speed		-1.35*10 ⁻⁸ *** (.0005)					
Sunshine - Change		.0196* (.0303)					
Interaction: SO ₂ * Diurnal temp.		.00323** (.0056)					
Interaction: Air pressure * Humidity				-.00000145** (.0049)			
Dust aerosols - Change						.00000104* (.0139)	
Dust aerosols - Squared						-1.42*10 ⁻¹² * (.0247)	

We also find clear differences between people with a high BMI compared to those with a low BMI. People with a high BMI are more likely to be hospitalised for pneumonia during times with high diurnal temperature, which is strengthened by high precipitation and increases in air pressure. At the same time, they seem to be less vulnerable to air pollution and the combination of heavy rainfall and sunshine within one day.

Especially interesting are the results for pneumonia patients with chronic pulmonary disease where we do not find any differences in the contemporary comparison and only two differences in the total comparison. Pneumonia appears more frequently among these people during times of high air pollution and low air pressure in combination with low humidity.

Additionally, while the triggering conditions (contemporary comparison) are not found to differ between the young and the elderly, we find that the relationship between pneumonia and minimum temperature in the total comparison is due to the older people (above 64 in age) in

the sample, while the younger people show no dependence. Older people show a strong relationship between being hospitalised with pneumonia and very low minimum temperature.

A comparison between female and male people only results in slightly significant (above 1% level) differences, indicating a lower reaction on the level of dust aerosols but higher reaction on changes of dust aerosols by females. Furthermore, pneumonia is more likely among females compared to males at times of low sunshine combined with low precipitation.

Moreover, we find a relationship between pneumonia and cardiac insufficiency. People with cardiac insufficiency are more likely to be hospitalised for pneumonia compared to other people during times with high diurnal temperature and low humidity. No differences are detected between pneumonia patients with and without diabetes regarding their sensitivity to meteorological and air conditions.

5. Conclusion

Our study adds to the current literature on the relationship of meteorological and air condition with pneumonia cases or deaths two perspectives, a long-term and a short-term perspective while at the same time account for diverse patients' personal and health background characteristics. The first perspective allows to distinguish between general conditions that are associated with higher frequencies of pneumonia cases whereas the second perspective focuses rather on triggering conditions that explain why pneumonia occurred at exactly this day and not a few days earlier or later. Triggering conditions are found to be especially strong during changes in meteorological and air conditions, namely decreases in sunshine hours and increases in diurnal temperature and air pollution as well as days with changes between sunshine and heavy rainfalls. Climate change is associated with more frequent, intense and variable meteorological conditions, so that we can expect higher numbers of pneumonia hospitalisations in the future.

The long-term analysis of the meteorological and air conditions co-occurring with higher frequencies of pneumonia confirms many of the findings in the literature. More specifically, we examine that temperature and air pollution seem to not directly influence pneumonia on their own but predominantly in combination with each other or with low air pressure. This implies that future research should avoid studying the impacts of meteorological and air conditions on pneumonia separately but in combination with each other. Additionally, more research should be conducted on specific meteorological constellations, meaning more complex interactions of meteorological and air conditions on pneumonia cases. Furthermore, future research should also pay attention to testing different lags since the selection of the best fitting lag strongly impacts the overall results as well as the suitability for different population groups.

The inclusion of various personal characteristics has shown that some findings do not hold for all people but only apply for certain subpopulations with a specific personal and/or health background. For example, changes in the diurnal temperature seem to trigger pneumonia only among people with a high BMI, decreases in sunshine hours seem to impact predominantly people with diabetes, and low temperatures significantly co-occur with pneumonia solely for the elderly. Many further relationships between personal characteristics and the relevance of certain meteorological and air conditions are found in this paper. Based on our findings, not

only resources and treatments at hospitals and doctor's offices during pneumonia-triggering meteorological and air conditions can be better prepared but also detailed and personal guiding prophylaxis can be advised to reduce the risks of the most vulnerable subgroups.

References

- Aliberti, S., Dela Cruz, C. S., Amati, F., Sotgiu, G., & Restrepo, M. I. (2021). Community-acquired pneumonia. *The Lancet*, 398(10303), 906–919. [https://doi.org/10.1016/S0140-6736\(21\)00630-9](https://doi.org/10.1016/S0140-6736(21)00630-9)
- Basu, R., & Samet, J. M. (2002). Relation between elevated ambient temperature and mortality: A review of the epidemiologic evidence. *Epidemiologic Reviews*, 24(2), 190–202. <https://doi.org/10.1093/epirev/mxf007>
- Behrendt, J., Penda, E., Finkler, A., Heil, U., & Polte-Rudolf, C. (2011). *Beschreibung der Datenbasis des NKDZ*. Offenbach.
- Benavides, R., Montes, F., Rubio, A., & Osoro, K. (2007). Geostatistical modelling of air temperature in a mountainous region of Northern Spain. *Agricultural and Forest Meteorology*, 146(3-4), 173–188. <https://doi.org/10.1016/j.agrformet.2007.05.014>
- Bianchi, E., Villalba, R., Viale, M., Couvreur, F., & Marticorena, R. (2016). New precipitation and temperature grids for northern Patagonia: Advances in relation to global climate grids. *Journal of Meteorological Research*, 30(1), 38–52. <https://doi.org/10.1007/s13351-015-5058-y>
- Bivand, R., Pebesma, E [EJ], & Gomez-Rubio, V. (2008). *Applied spatial data analysis with R*. <https://link.springer.com/content/pdf/10.1007/978-1-4614-7618-4.pdf>
- Bollmeyer, C., Keller, J. D., Ohlwein, C., Wahl, S., Crewell, S., Friederichs, P., Hense, A., Keune, J., Kneifel, S., Pschardt, I., Redl, S., & Steinke, S. (2015). Towards a high-resolution regional reanalysis for the European CORDEX domain. *Quarterly Journal of the Royal Meteorological Society*, 141(686), 1–15. <https://doi.org/10.1002/qj.2486>
- Braga, A. L. F., Zanobetti, A., & Schwartz, J [Joel] (2002). The effect of weather on respiratory and cardiovascular deaths in 12 U.S. Cities. *Environmental Health Perspectives*, 110(9), 859–863. <https://doi.org/10.1289/ehp.02110859>
- Brenner, T., & Pudelko, F. (2019). The effects of public research and subsidies on regional structural strength. *Journal of Evolutionary Economics*, 29(5), 1433–1458. <https://doi.org/10.1007/s00191-019-00626-x>
- Chan, P. K., Chew, F. T., Tan, T. N., Chua, K. B., & Hooi, P. S. (2002). Seasonal variation in respiratory syncytial virus chest infection in the tropics. *Pediatric Pulmonology*, 34(1), 47–51.
- Chiu, H.-F., Cheng, M.-H., & Yang, C.-Y. (2009). Air pollution and hospital admissions for pneumonia in a subtropical city: Taipei, Taiwan. *Inhalation Toxicology*, 21(1), 32–37. <https://doi.org/10.1080/08958370802441198>
- Chung, J.-Y., Honda, Y., Hong, Y.-C., Pan, X.-C., Guo, Y.-L., & Kim, H. (2009). Ambient temperature and mortality: An international study in four capital cities of East Asia. *The Science of the Total Environment*, 408(2), 390–396. <https://doi.org/10.1016/j.scitotenv.2009.09.009>
- Copernicus Climate Change Service. (2020). *Extreme precipitation risk indicators for Europe and European cities from 1950 to 2019*. <https://doi.org/10.24381/cds.3a9c4f89>
- Deal, E., McFadden, E., Ingram, R., Breslin, F., & Jaeger, J. (1980). Airway Responsiveness to Cold Air and Hyperpnea in Normal Subjects and in Those with Hay Fever and Asthma1–3. *American Review of Respiratory Disease*.
- DWD Climate Data Center. (2021). *Historische tägliche Stationsbeobachtungen (Temperatur, Druck, Niederschlag, Sonnenscheindauer, etc.) für Deutschland*.
- Eccles, R., & Wilkinson, J. E. (2015). Exposure to cold and acute upper respiratory tract infection. *Rhinology*, 53(2), 99–106. <https://doi.org/10.4193/Rhino14.239>
- Europäische Kommission - Eurostat/GISCO. (2022). *Administrative Units, Statitital Units (2016)*. 15.01.2022

- Falagas, M. E., Theocharis, G., Spanos, A., Vlara, L. A., Issaris, E. A., Panos, G., & Peppas, G. (2008). Effect of meteorological variables on the incidence of respiratory tract infections. *Respiratory Medicine*, 102(5), 733–737.
- Ge, W. Z., Xu, F., Zhao, Z. H., Zhao, J. Z., & Kan, H. D. (2013). Association between diurnal temperature range and respiratory tract infections. *Biomedical and Environmental Sciences: BES*, 26(3), 222–225. <https://doi.org/10.3967/0895-3988.2013.03.009>
- Gillissen, A., & Welte, T [Tobias]. (2014). *Herausforderungen, Zukunftsperspektiven, Forschungsansätze*. FRISCHTEXTE Verlag.
- Gräler, B., Pebesma, E., & Heuvelink, G. (2016). Spatio-Temporal Interpolation using gstat. *The R Journal*, 8(1), 204. <https://doi.org/10.32614/rj-2016-014>
- Green, R. S., Basu, R., Malig, B., Broadwin, R., Kim, J. J., & Ostro, B. (2010). The effect of temperature on hospital admissions in nine California counties. *International Journal of Public Health*, 55(2), 113–121. <https://doi.org/10.1007/s00038-009-0076-0>
- Höffken, G., Lorenz, J., Kern, W., Welte, T [T.], Bauer, T., Dalhoff, K [K.], Dietrich, E., Ewig, S., Gastmeier, P., Grabein, B., Halle, E., Kolditz, M., Marre, R., & Sitter, H. (2009). Epidemiologie, Diagnostik, antimikrobielle Therapie und Management von erwachsenen Patienten mit ambulant erworbenen unteren Atemwegsinfektionen sowie ambulant erworbener Pneumonie--Update 2009. S3-Leitlinie der Paul-Ehrlich-Gesellschaft für Chemotherapie, der Deutschen Gesellschaft für Pneumologie und Beatmungsmedizin, der Deutschen Gesellschaft für Infektiologie und vom Kompetenznetzwerk CAPNETZ [Epidemiology, diagnosis, antimicrobial therapy and management of community-acquired pneumonia and lower respiratory tract infections in adults. Guidelines of the Paul-Ehrlich-Society for Chemotherapy, the German Respiratory Society, the German Society for Infectiology and the Competence Network CAPNETZ Germany]. *Pneumologie*, 63(10), e1-68. <https://doi.org/10.1055/s-0029-1215037>
- Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., & Suttie, M. (2019). The CAMS reanalysis of atmospheric composition. *Atmospheric Chemistry and Physics*, 19(6), 3515–3556. <https://doi.org/10.5194/acp-19-3515-2019>
- Institut für Qualitätssicherung und Transparenz im Gesundheitswesen. (2016). *Bundesauswertung zum Erfassungsjahr 2015. Ambulant erworbene Pneumonie. Qualitätsindikatoren*. <http://docplayer.org/64901719-Bundesauswertung-zum-erfassungsjahr-ambulant-erworbene-pneumonie-qualitaetsindikatoren.html>
- IPCC. (2022). *Impacts, Adaptation and Vulnerability. Working Group II Contribution to the IPCC Sixth Assessment Report*. Cambridge University Press. <https://www.ipcc.ch/assessment-report/ar6/>
- Jarvis, A., Reuter, H., Nelson, A., & Guevara, E. (2008). *Hole-filled seamless SRTM data V4*, International Centre for Tropical Agriculture CIAT., available from <https://srtm.csi.cgiar.org> <<https://srtm.csi.cgiar.org/>>. <https://scholar.google.com/citations?user=6lajxpkaaaaj&hl=de&oi=sra>
- Kaspar, F., Müller-Westermeier, G., Penda, E., Mächel, H., Zimmermann, K., Kaiser-Weiss, A., & Deutschländer, T. (2013). Monitoring of climate change in Germany – data, products and services of Germany's National Climate Data Centre. In *Advances in Science and Research* (pp. 99–106). Copernicus GmbH. <https://doi.org/10.5194/asr-10-99-2013>
- Katsouyanni, K., Touloumi, G., Spix, C., Schwartz, J [J.], Balducci, F., Medina, S., Rossi, G., Wojtyniak, B., Sunyer, J., Bacharova, L., Schouten, J. P., Ponka, A., & Anderson, H. R. (1997). Short-term effects of ambient sulphur dioxide and particulate

- matter on mortality in 12 European cities: Results from time series data from the APHEA project. *Air Pollution and Health: A European Approach. BMJ (Clinical Research Ed.)*, 314(7095), 1658–1663. <https://doi.org/10.1136/bmj.314.7095.1658>
- Kim, J., Kim, J.-H., Cheong, H.-K., Kim, H., Honda, Y., Ha, M., Hashizume, M [Masahiro], Kolam, J., & Inape, K. (2016). Effect of Climate Factors on the Childhood Pneumonia in Papua New Guinea: A Time-Series Analysis. *International Journal of Environmental Research and Public Health*, 13(2), 213.
- Kwak, H.-Y., Ko, J., Lee, S., & Joh, C.-H. (2017). Identifying the correlation between rainfall, traffic flow performance and air pollution concentration in Seoul using a path analysis. *Transportation Research Procedia*, 25, 3552–3563. <https://doi.org/10.1016/j.trpro.2017.05.288>
- Lam, H. C.-Y., Chan, E. Y.-Y., & Goggins, W. B. (2018). Comparison of short-term associations with meteorological variables between COPD and pneumonia hospitalization among the elderly in Hong Kong-a time-series study. *International Journal of Biometeorology*, 62(8), 1447–1460. <https://doi.org/10.1007/s00484-018-1542-2>
- Larsson, K., Ohlsén, P., Larsson, L., Malmberg, P., Rydström, P. O., & Ulriksen, H. (1993). High prevalence of asthma in cross country skiers. *BMJ (Clinical Research Ed.)*, 307(6915), 1326–1329. <https://doi.org/10.1136/bmj.307.6915.1326>
- Larsson, K., Tornling, G., Gavhed, D., Müller-Suur, C., & Palmberg, L. (1998). Inhalation of cold air increases the number of inflammatory cells in the lungs in healthy subjects. *The European Respiratory Journal*, 12(4), 825–830. <https://doi.org/10.1183/09031936.98.12040825>
- Lim, Y.-H., Hong, Y.-C., & Kim, H. (2012). Effects of diurnal temperature range on cardiovascular and respiratory hospital admissions in Korea. *The Science of the Total Environment*, 417-418, 55–60. <https://doi.org/10.1016/j.scitotenv.2011.12.048>
- Liu, L., Breitner, S., Pan, X., Franck, U., Leitte, A. M., Wiedensohler, A., Klot, S. von, Wichmann, H.-E., Peters, A., & Schneider, A. (2011). Associations between air temperature and cardio-respiratory mortality in the urban area of Beijing, China: A time-series analysis. *Environmental Health: A Global Access Science Source*, 10, 51.
- Liu, Y., Kan, H., Xu, J [Jianming], Rogers, D., Peng, L., Ye, X., Chen, R., Zhang, Y., & Wang, W. (2014). Temporal relationship between hospital admissions for pneumonia and weather conditions in Shanghai, China: A time-series analysis. *BMJ Open*, 4(7), e004961.
- Mäkinen, T. M., Juvonen, R., Jokelainen, J., Harju, T. H., Peitso, A., Bloigu, A., Silvennoinen-Kassinen, S., Leinonen, M., & Hassi, J. (2009). Cold temperature and low humidity are associated with increased occurrence of respiratory tract infections. *Respiratory Medicine*, 103(3), 456–462. <https://doi.org/10.1016/j.rmed.2008.09.011>
- Murdoch, D. R., & Jennings, L. C. (2009). Association of respiratory virus activity and environmental factors with the incidence of invasive pneumococcal disease. *The Journal of Infection*, 58(1), 37–46. <https://doi.org/10.1016/j.jinf.2008.10.011>
- Nascimento-Carvalho, C. M., Cardoso, M.-R. A., Barral, A., Araújo-Neto, C. A., Oliveira, J. R., Sobral, L. S., Saukkoripi, A., Paldanius, M., Vainionpää, R., Leinonen, M., & Ruuskanen, O. (2010). Seasonal patterns of viral and bacterial infections among children hospitalized with community-acquired pneumonia in a tropical region. *Scandinavian Journal of Infectious Diseases*, 42(11-12), 839–844. <https://doi.org/10.3109/00365548.2010.498020>
- Nhung, N. T. T., Amini, H., Schindler, C., Kutlar Joss, M., Dien, T. M., Probst-Hensch, N., Perez, L., & Künzli, N. (2017). Short-term association between ambient air pollution and pneumonia in children: A systematic review and meta-analysis of time-series and

- case-crossover studies. *Environmental Pollution (Barking, Essex: 1987)*, 230, 1000–1008. <https://doi.org/10.1016/j.envpol.2017.07.063>
- Onozuka, D., Hashizume, M [M.], & Hagihara, A. (2009). Impact of weather factors on *Mycoplasma pneumoniae pneumonia*. *Thorax*, 64(6), 507–511. <https://doi.org/10.1136/thx.2008.111237>
- Paynter, S., Weinstein, P., Ware, R. S., Lucero, M. G., Tallo, V., Nohynek, H., Barnett, A. G., Skelly, C., Simoes, E. A., Sly, P. D., & Williams, G. (2013). Sunshine, rainfall, humidity and child pneumonia in the tropics: Time-series analyses. *Epidemiology and Infection*, 141(6), 1328–1336. <https://doi.org/10.1017/s0950268812001379>
- Pebesma, E. J. (2004). Multivariable geostatistics in S: the gstat package. *Computers & Geosciences*, 30(7), 683–691. <https://doi.org/10.1016/j.cageo.2004.03.012>
- Plass, D., Vos, T., Hornberg, C., Scheidt-Nave, C., Zeeb, H., & Krämer, A. (2014). *Entwicklung der Krankheitslast in Deutschland*. Robert Koch-Institut, Epidemiologie und Gesundheitsberichterstattung. <https://edoc.rki.de/handle/176904/1981> <https://doi.org/10.25646/1906>
- R Core Team. (2021). *R: A language and environment for statistical computing*. <http://r.meteo.uni.wroc.pl/web/packages/dplr/vignettes/intro-dplr.pdf>
- Reuter, H. I., Nelson, A., & Jarvis, A. (2007). An evaluation of void-filling interpolation methods for SRTM data. *International Journal of Geographical Information Science*, 21(9), 983–1008. <https://doi.org/10.1080/13658810601169899>
- RKI. (2017). *Zeitliche Trends in der Inzidenz und Sterblichkeit respiratorischer Krankheiten von hoher Public-Health-Relevanz in Deutschland*. RKI-Bib1 (Robert Koch-Institut). <https://edoc.rki.de/handle/176904/2780> <https://doi.org/10.17886/RKI-GBE-2017-049>
- Samet, J., Zeger, S., Kelsall, J., Xu, J [J.], & Kalkstein, L. (1998). Does weather confound or modify the association of particulate air pollution with mortality? An analysis of the Philadelphia data, 1973–1980. *Environmental Research*, 77(1), 9–19. <https://doi.org/10.1006/enrs.1997.3821>
- Schnoor, M., Hedicke, J., Dalhoff, K [Klaus], Raspe, H., & Schäfer, T. (2007). Approaches to estimate the population-based incidence of community acquired pneumonia. *The Journal of Infection*, 55(3), 233–239. <https://doi.org/10.1016/j.jinf.2007.04.355>
- Sohn, S., Cho, W., Kim, J. A., Altaluoni, A., Hong, K., & Chun, B. C. (2019). 'pneumonia Weather': Short-term Effects of Meteorological Factors on Emergency Room Visits Due to Pneumonia in Seoul, Korea. *Journal of Preventive Medicine and Public Health = Yebang Uihakhoe Chi*, 52(2), 82–91.
- Wang, Z.-B., Ren, L., Lu, Q.-B., Zhang, X.-A., Miao, D., Hu, Y.-Y., Dai, K., Li, H., Luo, Z.-X., Fang, L.-Q., Liu, E.-M., & Liu, W. (2021). The Impact of Weather and Air Pollution on Viral Infection and Disease Outcome Among Pediatric Pneumonia Patients in Chongqing, China, from 2009 to 2018: A Prospective Observational Study. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*, 73(2), e513–e522. <https://doi.org/10.1093/cid/ciaa997>
- World Health Organization. (2007). *Global Surveillance, Prevention and Control of Chronic Respiratory Diseases: A Comprehensive Approach*. World Health Organization.
- Xu, Z., Huang, C., Su, H., Turner, L. R., Qiao, Z., & Tong, S. (2013). Diurnal temperature range and childhood asthma: A time-series study. *Environmental Health: A Global Access Science Source*, 12, 12.

Appendix A




Figure A2: Pneumonia Deaths per 100,000 inhabitants (age-standardized): Germany.




Figure A3: Number of people hospitalized for pneumonia: Germany.