




MONITORING SOIL PARAMETERS: EARTH OBSERVATION AS A TRUSTWORTHY TECHNIQUE

Earth Observation as a trustworthy technique for monitoring soil parameters
(6 minute presentations)

*Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra*



ESA Symposium on Earth Observation for Soil Protection and Restoration



Our soils, the earth upper surface, with its key role for terrestrial life functions: plant growth, gas exchange with the atmosphere, retention of water and nutrients, among the objectives of many of us, ...Soils are being rediscovered from numerous stand viewpoints: biochemical, ecologic, technical, analytical, social, policy, economic, etc..


Earth Observation (EO) and Copernicus data are needed for achieving the objectives, the scientific community, “policy stakeholders” will serve to monitor the requirements of the various soil health indicators stated in the law



Joint Research Centre




1. We need data flows to populate indicators of soil health and create knowledge for policy



2. **EUSO Soil Dashboard** is based on the convergence of scientific evidence – question how to monitor change. How can EO help?

Convergence of scientific evidence

- 18 Soil degradation indicators
- 63 % of unhealthy soils
- Dashboard shows location and different types of soil degradation in the EU




ESA Symposium on Earth Observation for Soil Protection and Restoration, 06 – 07 March 2024, ESA-ESRIN, Frascati (Rome), Italy

Assessing Policy Impact

<https://esdac.jrc.ec.europa.eu/esdacviewer/euso-dashboard/>




3. Out of all indicators, only soil sealing comes from EO directly, in other indicators EO data are co-variant




Soil degradation indicators

Select a soil degradation indicator:



Indicator Nuts 0 Nuts 2



DAY 1 Panelist presentation SESSION 2 10:00 MONITORING SOIL PARAMETERS: EARTH OBSERVATION AS A TRUSTWORTHY TECHNIQUE

Earth Observation as a trustworthy technique for monitoring soil parameters

Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra

4. Imperviousness/ Sealing Copernicus Land Monitoring Service CLMS



[CLMS portfolio](#) [Dataset catalogue](#) [Data viewer](#) [Use cases](#) [About us](#)


[Home](#) > [CLMS portfolio](#) > [High Resolution Layer Impervious Built-up](#) > [Impervious Built-up 2018 \(raster 10 m and 100 m\), Europe, 3-yearly](#)

Impervious Built-up 2018 (raster 10 m and 100 m), Europe, 3-yearly

General info

Provides at pan-European level in the spatial resolution of 10 m binary information (building / no building) within the sealing outline derived from the Imperviousness Density 2018 including the aggregated Share of Built-up layer in the spatial resolution of 100 m showing the percentage of built-up areas across Europe.

[Download](#)



[View in the data viewer](#)

Validation status

Validated

Dataset citation

- DOI (raster 100 m): <https://doi.org/10.2909/a807e528-431a-4dca-a6cd-0e8947563fce>
- DOI (raster 10 m): <https://doi.org/10.2909/3e412def-a4e6-4413-98bb-42b571afd15e>

Services

[WMS](#)

[REST API](#)

You can find instructions on how to cite CLMS data in our [Data policy](#) section.

Characteristics

Release / Major version:
V1

Temporal extent:
2017-2019

Type:
Satellite products

Position accuracy:
According to orthorectified satellite image

Projection:
EPSG:3035

Temporal usability:
Archive with regular updates

Platform:
Sentinel-2


Thematic accuracy:
≥ 90%

Spatial coverage:
Europe

Update frequency:
3-yearly

Spatial resolution:

Datasets




Imperviousness Density 2018 (raster 10 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 10 m and 100 m the sealing density in the range from 0% to 100% for the 2018 reference year.

[View more](#)

[Download](#)

[View in the data viewer](#)




Imperviousness Density 2015 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2015 reference year.

[View more](#)

[Download](#)

[View in the data viewer](#)




Imperviousness Density 2012 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2012 reference year.

[View more](#)

[Download](#)

[View in the data viewer](#)



Imperviousness Density 2009 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2009 reference year.

[View more](#)

[Download](#)

[View in the data viewer](#)



Imperviousness Density 2006 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2006 reference year.

5. Other indicators currently possible from EO ?

→ EO data for **Management Practices** (tillage intensity, cover crops, irrigation)

→ EO data used as parameters of models or as covariates for **soil spatial modelling** (geostatistics and AI)

→ EO data high resolution **Digital Elevation Models** and **Vegetation indices** (Net primary productivity, biomass)

→ EO data for **soil salinity**

6. Some 1st principles, to classify and monitor land objects with EO we are trying to use reflected energy, radar frequencies

→ this implies that by using EO as a proxy for soil assessment we replicate complex laboratory techniques e.g.

- Nutrient content
- N
- P
- K
- Micronutrients (Zn, Br, Mn, ...)
- Abundance of salts
- NaCl
- K₂O
- Texture,
- Clay
- Sand
- Silt
- Moisture
- Wilting point
- Ksat
- Temperature
- T_{max}
- T_{min}
- Prolonged droughts

7. Many of promising case studies on soils (you can find some example, soil moisture, albedo) but to date, very limited operational services specifically on soils.



Burnt Area	NDVI
Dry Matter Prod.	Soil Water Index
FAPAR	Surf. Soil Moisture
FCOVER	VCI
Leaf Area Index	VPI
Land Cover	

Normalized Difference Vegetation Index

The Normalized Difference Vegetation Index (NDVI) is an indicator of the greenness of the biomes. Even though it is not a physical property of the vegetation cover, its very simple formulation $NDVI = (REF_{nir} - REF_{red}) / (REF_{nir} + REF_{red})$ where REF_{nir} and REF_{red} are the spectral reflectances measured in the near infrared and red wavebands respectively, makes it widely used for ecosystems monitoring.

NDVI product updates

NDVI 300m - reprocessing for missing blocks in October and November 2023 products

Fri, 22 Dec 2023

NDVI 300m for July 2022 reprocessed

Mon, 27 Mar 2023


Often these products comes in temporal series and need to be harmonized to be effective for soil modelling

8. Now we have the WORLDSOILS Carbon product

→ It need commitment from ESA/COPERNICUS for repeat and continued development



→ Continuous development and **temporal release** of the product will allow for wide usage and extensive validation (e.g with LUCAS soil organic carbon).




DAY 1 Panelist presentation SESSION 2 10:00 MONITORING SOIL PARAMETERS: EARTH OBSERVATION AS A TRUSTWORTHY TECHNIQUE

Earth Observation as a trustworthy technique for monitoring soil parameters

Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra

9. Lot of promise in COPERNICUS Expansion Programme this will be discussed in the session 5





DAY 1 Chair panel 15:45

SOC Maps in Practical Use. Chairs:
Calogero Schillaci (EUSO-JRC), Bas
van Wesemael (UCLouvain)


PANEL: SOC Maps relevance
and usability for the
institutional mandate



ESA Symposium on Earth Observation for Soil Protection and Restoration

JRC has developed several SOC assessment at European Scale based on LUCAS


- These are used in policy implementation and monitoring (e.g CAP, SML, NRL, CRCF)
- Thanks to the unprecedented EO high resolution (spectral, spatial and temporal) over continental areas, are an essential component for monitoring the SOC seamlessly



A map of the topsoil organic carbon content of Europe generated by a generalized additive model, De Brogniez et al 2015



Assessment of soil organic carbon stocks under future climate and land cover changes in Europe, Yigini and Panagos 2016,



Soil organic carbon stocks in European croplands and grasslands: How much have we lost in the past decade? De Rosa et al. 2024



DAY 1 WORLDSOILS SOC Prediction Maps Results and Way Forward 16:30-17.



Conclusion and way forward

ESA Symposium on Earth Observation for Soil Protection and Restoration



Day 2 9:00-9:30 **PARALLEL SESSIONS IN
MAGELLAN (Hall 2)**

**Keynote SESSION 5 - ADVANCED SATELLITE
INSTRUMENTATION FOR SOIL HEALTH**

*Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra*



ESA Symposium on Earth Observation for Soil Protection and Restoration


Day 2 9:00-9:30 PARALLEL SESSIONS IN MAGELLAN (Hall 2)

Keynote SESSION 5 - ADVANCED SATELLITE INSTRUMENTATION FOR SOIL HEALTH


Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra

1. Why we need EO for soil monitoring?

- Monitoring our soils
- Earth Observation (EO) and Copernicus
- The scientific community
- Policy stakeholders
- Soil health indicators stated in the law



Joint
Research
Centre



Day 2 9:00-9:30 PARALLEL SESSIONS IN MAGELLAN (Hall 2)

Keynote SESSION 5 - ADVANCED SATELLITE INSTRUMENTATION FOR SOIL HEALTH

Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra

2. Scientific peer-review literature

European Space Agency - ESA

8-10 rue Mario Nikis, Paris
France
Affiliation ID: 60029398
Other name formats: [European Space Agency](#) [European Space Agency \(esa\)](#) [Esa](#) [European Space Agency \(esa-ested\)](#) [European Space Agency \(esa/ested\)](#)
[European Space Agency \(esa\)](#) [Esa Neo Coordination Centre](#) [Rhea For European Space Agency \(esa\)](#)
[Aurora Technology For European Space Agency \(esa\)](#) [Esa Headquarters](#)

Affiliation profile actions


- [Give feedback](#)
- [Set document alert](#)
- [Export subject area data](#)

Documents by subject area Collaborating affiliations Documents by source

Sort by: [Document count \(high-low\)](#)

Subject Area	Document Count	Subject Area	Document Count
Earth and Planetary Sciences	4101	Energy	89
Engineering	3258	Pharmacology, Toxicology and Pharmaceuticals	68
Physics and Astronomy	3027	Neuroscience	44
Computer Science	1264	Arts and Humanities	36
Materials Science	477	Economics, Econometrics and Finance	36
Mathematics	364	Decision Sciences	32
Social Sciences	313	Immunology and Microbiology	30
Biochemistry, Genetics and Molecular Biology	296	Business, Management and Accounting	25
Agricultural and Biological Sciences	292	Nursing	18
Environmental Science	291	Health Professions	15
Chemistry	277	Psychology	8
Medicine	249	Veterinary	2
Multidisciplinary	103	Dentistry	1

European Space Agency - ESA



ESRIN - ESA Centre for Earth Observation

Frascati
RM, Italy
Affiliation ID: 60073749
Other name formats: [Esrin](#) [Esa-esrin](#) [Esa/esrin](#) [Esa Esrin](#) [European Space Research Institute](#) [European Space Research Institute \(esrin\)](#)
[European Space Agency](#) [C/o Esrin](#) [European Space Agency \(esa-esrin\)](#) [European Space Agency/esrin](#)

Affiliation profile actions


- [Give feedback](#)
- [Set document alert](#)
- [Export subject area data](#)

Documents by subject area Collaborating affiliations Documents by source

Sort by: [Document count \(high-low\)](#)

Subject Area	Document Count	Subject Area	Document Count
Earth and Planetary Sciences	1504	Multidisciplinary	23
Engineering	759	Chemical Engineering	20
Physics and Astronomy	546	Chemistry	20
Computer Science	420	Decision Sciences	11
Environmental Science	143	Medicine	10
Agricultural and Biological Sciences	109	Biochemistry, Genetics and Molecular Biol...	8
Social Sciences	79	Business, Management and Accounting	5
Mathematics	68	Economics, Econometrics and Finance	2
Materials Science	53	Arts and Humanities	1
Energy	52	Neuroscience	1

ESRIN - ESA Centre for Earth Observation



Day 2 9:00-9:30 PARALLEL SESSIONS IN MAGELLAN (Hall 2)

Keynote SESSION 5 - ADVANCED SATELLITE INSTRUMENTATION FOR SOIL HEALTH

Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra

2. Scientific peer-review literature

The screenshot shows a Scopus search results page. At the top, the Scopus logo is on the left, and navigation links for Search, Lists, Sources, SciVal, and user options (Create account, Sign in) are on the right. The search results section displays "59 document results" for the keyword "SOIL", which is highlighted with a blue arrow. Below this, a search query is shown in a red box: `(AF-ID ("European Space Agency - ESA" 60029398) AND SUBJAREA (agri)) AND ("soil")`. The results are sorted by "Date (newest)". A table lists the top three results, including document titles, authors, years, sources, and citation counts.

	Document title	Authors	Year	Source	Cited by
<input type="checkbox"/>	1 Space hardware for concrete sample production on ISS "MASON concrete mixer" <i>Open Access</i>	Müller, J.T.I., Rattenbacher, B., Tell, K., (...), Sperl, M., Schnellenbach-Held, M.	2023	npj Microgravity 9(1),57	0
<input type="checkbox"/>	2 The EnMAP imaging spectroscopy mission towards operations <i>Open Access</i>	Storch, T., Honold, H.-P., Chabrilat, S., (...), Schickling, A., Fischer, S.	2023	Remote Sensing of Environment 294,113632	14
<input type="checkbox"/>	3 Constraining industrial ammonia emissions using hyperspectral infrared imaging	Noppen, L., Clarisse, L., Tack, F., (...), Schuettemeyer, D., Coheur	2023	Remote Sensing of Environment	0

Day 2 9:00-9:30 PARALLEL SESSIONS IN MAGELLAN (Hall 2)

Keynote SESSION 5 - ADVANCED SATELLITE INSTRUMENTATION FOR SOIL HEALTH

Calogero Schillaci and the EUSO Team
European Soil Observatory
European Commission - JRC Ispra

2. Scientific peer-review literature

Brought to you by EC - Joint Research Centre



Search Lists Sources SciVal ? Sign in Create account

Welcome to a more intuitive and efficient search experience. [See what is new](#)

Advanced query

Filters [Clear all](#)

Year [Clear](#)

Range Individual



2017 - 2024

229 documents found

	Document title	Authors	Source	Year	Citations
<input type="checkbox"/> 1	Article • Open access A comparative analysis of machine learning techniques for aboveground biomass estimation: A case study of the Western Ghats, India	Ayushi, K., Babu, K.N., Ayyappan, N., ...Kakkara, A., Reddy, C.S.	Ecological Informatics, 80, 102479	2024	0
	Show abstract View at Publisher Related documents				
<input type="checkbox"/> 2	Article Spatial and temporal variation of soil properties and soil organic carbon in semi-arid areas of Sub-Saharan Africa	Suh, C.N., Tsheko, R.	Geoderma Regional, 36, e00770	2024	1
	Show abstract View at Publisher Related documents				

3. To populate the **EUSO dashboard** with soil indicators we need numerous **EO data** to provide valuable indicators of soil health

Soil degradation indicators

- Copper: copper contamination
- Mercury: mercury contamination
- Zinc: zinc contamination
- Compac.: soil compaction
- Erosion - till: tillage erosion
- Erosion - wa: water erosion
- Erosion - wi: wind erosion
- N surplus: nitrogen surplus
- P deficit: phosphorus deficiency
- P excess: phosphorus excess
- Peatland degr.: loss of organic soils
- Erosion - fire: fire erosion
- Erosion - har: harvest erosion
- Saliniz.: soil salinization
- Soil bio.: loss of soil biodiversity
- SOC: loss of soil organic carbon


Soil degradation indicators

Select a soil degradation indicator:

3. EUSO Soil Dashboard is based on the convergence of scientific evidence – question how to monitor change. How can EO help?

Convergence of scientific evidence

- 18 Soil degradation indicators
- 63 % of unhealthy soils
- Dashboard shows location and different types of soil degradation in the EU



Proportion of land affected by soil degradation in the EU

Assessing Policy Impact

<https://esdac.jrc.ec.europa.eu/esdacviewer/euso-dashboard/>



3. In EUSO Dashboard **Convergence of Evidence** is used to stack together the soil threats and determine how much pressure is occurring in each unit (pixel)

→ question **how to monitor change**. How can EO help map soil degradation processes?

As a commitment on the soil strategy the soil monitoring law is expected to come bringing some key changes in EU

- By **2050**, all EU soil ecosystems are in **healthy** condition and are thus more **resilient**, which will require very **decisive changes** in this decade.
- By then, **protection, sustainable use and restoration of soil** has become the norm.
- Healthy soils are essential to achieve **climate neutrality**, a clean and **circular economy**, revert **biodiversity loss**, safeguard **human health**, halt **desertification** and revert **land degradation**.
- Legislative proposal by 2023

4. The new soil monitoring directive¹ relies on soil monitoring network and soil Indicators – are we able to derive them from EO – why not?
 → **Sensors** and products predominantly designed for vegetation.
 → In addition, remote sensing data **acquisition** and **accuracy** can be limited under dense vegetation cover with the obstructed view of the **soil surface** making it impossible to directly detect soil characteristics.

Soil descriptor	aspect of soil degradation
Electrical Conductivity	salinization
Soil erosion rate	loss of soil by erosion
SOC concentration	loss of organic matter
Bulk density in subsoil	compaction
Bulk density in topsoil	
Extractable phosphorus	excess nutrients
Nitrogen in soil	
Concentration of heavy metals	contamination
Concentration of other contaminants	
Soil water holding capacity	loss of capacity for water retention
Soil acidity (pH)	acidification
Soil basal respiration	loss of soil biodiversity

Soil basal respiration	loss of soil biodiversity
------------------------	---------------------------

5. Trying to use **measurement of reflectance** to replicate **complex laboratory techniques**.
 EO data are proxies.

→ Refers to indirect measurements used to estimate **plant or vegetation status**, or microwave data can estimate **soil moisture content**. (E.g. LAI, measurements obtained by Neural network algorithm) provide valuable information for soil monitoring, but they often require **validation** against ground-based measurements for accuracy

Leaf Area Index

The Leaf Area Index is defined as half the total area of green elements of the canopy per unit horizontal ground area. The satellite-derived value corresponds to the total green LAI of all the canopy layers, including the understory which may represent a very significant contribution, particularly for forests. Practically, the LAI quantifies the thickness of the vegetation cover.

LAI is recognized as an Essential Climate Variable (ECV) by the Global Climate Observing System (GCOS).

LAI product updates

- Geolocation correction in Sentinel-3-based vegetation products
 Fri, 19 Aug 2022
 - Geolocation issue in Sentinel-3-based vegetation products
 Thu, 12 May 2022
 - Sentinel 3 LAI, FAPAR and FCOVER product quality artifacts identified
 Thu, 07 Apr 2022
- [Read more or Subscribe](#)

LAI characteristics


LAI 300m
LAI 1km

Access
Algorithm
Quality
Application
Technical
Documents
Gallery

The details of the algorithm can be found in the ATBD.

Version	Main elements	Differences with previous version	Reference
1	<ul style="list-style-type: none"> Daily LAI 300m is estimated by applying a Neural Network on <ul style="list-style-type: none"> instantaneous Top-of-Canopy reflectances from Sentinel-3 OLCI (v1.1 products), or daily Top-of-Aerosol input reflectances from PROBA-V (v1.0). Temporal smoothing and small gap filling is applied to the instantaneous LAI estimates, discriminating Evergreen Broadleaf Forest (EBF) and no-EBF pixels. Temporal compositing is adapted to provide a near-real time (10-daily) estimate and successive updated estimates until a consolidated value is reached after about 2 months. 	Differences with 1km products: <ul style="list-style-type: none"> Input reflectances are corrected from ozone, water vapour, and surface pressure. Shortwave infrared reflectances and climatology information are omitted. Discrimination of EBF and no-EBF pixels. Algorithm runs on 300m PROBA-V only. 	

6. However, clear that **traditional soil mapping** involves labour-intensive field surveys, which can be expensive and time-consuming so we should be able to better benefit from **EO data** using space-borne sensor that provide a **synoptic view of the soils and its properties dynamics**.



Soil Map of the European Union and bordering countries (source: Soil Atlas of Europe).

European soil data Centre ESDAC


7. Need targeted sensors/instrumentation – **Soil Indices**

→ Integrated sensor products (e.g. thermal-SAR-Optical)

→ Example of LIDAR + vegetation for land degradation

Slope+ vegetation (biomass, NPP, NDVI or LAI)

→ Include the Sentinel 5 in the soil monitoring tools, Methane (CH₄), as the second most important GHG following carbon dioxide, holds significant implications for global warming and the carbon cycle (IPCC, [2021](#)).



The new relevant sensors in the Copernicus Extension Programme can provide for soils and in particular the SML and other policy areas (e.g. pollution, desertification)


8. Altimetry LIDAR and soils modelling, we need better Digital Elevation Models for:

→ Mapping pedogenic processes,


- soil formation (possible to some extent),
- horizonation (challenging but feasible with the radar application for the topsoil)

→ Monitoring erosion:

- **Water erosion** (sheet, rill, gully and Badland)
- **Wind erosion** (sand dunes, sand storms)
- Coastal erosion (riverbank erosion and coastline retreat)




Panagos et al., 2015 The new assessment of soil loss by water erosion in Europe



Borrelli et al, 2017 A New Assessment of Soil Loss Due to Wind

9. Unmanned aerial vehicles “Drones” are increasingly used for soil mapping?

→ Drones particularly useful for localized, high-resolution mapping and monitoring tasks



100m

→ EO via RS offers a wide coverage area, suitable for large-scale regional mapping and monitoring, can be acquired at regular intervals, cost-effective

10

Imperviousness/ Sealing Copernicus Land Monitoring Service CLMS



[CLMS portfolio](#) [Dataset catalogue](#) [Data viewer](#) [Use cases](#) [About us](#)


[Home](#) > [CLMS portfolio](#) > [High Resolution Layer Impervious Built-up](#) > [Impervious Built-up 2018 \(raster 10 m and 100 m\), Europe, 3-yearly](#)

Impervious Built-up 2018 (raster 10 m and 100 m), Europe, 3-yearly

General info

Provides at pan-European level in the spatial resolution of 10 m binary information (building / no building) within the sealing outline derived from the Imperviousness Density 2018 including the aggregated Share of Built-up layer in the spatial resolution of 100 m showing the percentage of built-up areas across Europe.

[Download](#)



[View in the data viewer](#)

Validation status

Validated

Dataset citation

- DOI (raster 100 m): <https://doi.org/10.2909/a807e528-431a-4dca-a6cd-0e8947563fce>
- DOI (raster 10 m): <https://doi.org/10.2909/3e412def-a4e6-4413-98bb-42b571afd15e>

Services

[WMS](#)

[REST API](#)

You can find instructions on how to cite CLMS data in our [Data policy](#) section.

Characteristics

Release / Major version:
V1

Temporal extent:
2017-2019

Type:
Satellite products

Position accuracy:
According to orthorectified satellite image

Projection:
EPSG:3035

Temporal usability:
Archive with regular updates

Platform:
Sentinel-2


Thematic accuracy:
≥ 90%

Spatial coverage:
Europe

Update frequency:
3-yearly

Spatial resolution:


Datasets



Imperviousness Density 2018 (raster 10 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 10 m and 100 m the sealing density in the range from 0% to 100% for the 2018 reference year.


[View more](#) [Download](#) [View in the data viewer](#)



Imperviousness Density 2015 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2015 reference year.


[View more](#) [Download](#) [View in the data viewer](#)



Imperviousness Density 2012 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2012 reference year.

[View more](#) [Download](#) [View in the data viewer](#)



Imperviousness Density 2009 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2009 reference year.

[View more](#) [Download](#) [View in the data viewer](#)



Imperviousness Density 2006 (raster 20 m and 100 m), Europe, 3-yearly

Provides at pan-European level in the spatial resolution of 20 m and 100 m the sealing density in the range from 0% to 100% for the 2006 reference year.


10. Also need for soil dedicated **sealing** detection and **classification** CLMS –



Even attempt to mitigate soil sealing not working

11 High-resolution (spatial and temporal) satellite imagery can be used as covariates of information in digital soil mapping


- Land surface temperature
- topography and derivatives (Slope, aspect, curvature, SPI, TWI, etc)
- Air fluxes/emissions mapping (CO₂, CH₄, NOX)
- Photogrammetry
- ESA's soil moisture



12 Soil Agricultural Practices

→ BARE SOIL there are several studies, but in EU it's usually due tillage and happen often in fall-winter cloudy months so application of optical remote sensing need to be complemented with the radar interferometry measurement.


→ AGRICULTURAL PRACTICES have been object of study by colleagues at the JRC



Matthews, F., et al., 2023. ISWCR, A field parcel-oriented approach to evaluate the crop cover-management factor and time-distributed erosion risk in Europe. Int. 10.1016/j.iswcr.2022.09.005

A.N. Fendrich et al.

Science of the Total Environment 873 (2023) 162300



Fendrich et al 2023 STOTEN From regional to parcel scale: A high-resolution map of cover crops across Europe combining satellite data with statistical surveys, [10.1016/j.scitotenv.2023.162300](https://doi.org/10.1016/j.scitotenv.2023.162300)

13. Hyperspectral

→ Enmap The Environmental Mapping and Analysis Program (*EnMAP*) is a German hyperspectral satellite mission that aims at monitoring and characterizing the Earth's measures geochemical, biochemical and biophysical variables providing information on the status and evolution of terrestrial and aquatic ecosystems

→ ASI "PRecursores IperSpettrale della Missione Applicativa" PRISMA lunched in 2019 in Kourou provides VNIR (Visible and Near-InfraRed), SWIR (Short-Wave InfraRed), at 30 m spatial resolution, and pancromatic at 5 m.



DAY 2 Closure 16.30-17:30 JRC