Journal Hopping – when journal de-indexing is not enough

Ambiguity

Ambiguity

Journal Hopping

ZIC2



User Guide

Save

Email

Advanced Create alert Create RSS

Send to

Sort by:

Most recent

♦ ↓<u>-</u>

Display options 🗱

TEXT AVAILABILITY

Abstract

Free full text

Full text



Save

Email

Send to

Sort by:

Most recent



Display options 🗱

TEXT AVAILABILITY

Abstract

Free full text

Full text





Search

User Guide

Save

Email

Send to

Sort by:

Most recent




Page

Display options 🗱

MY NCBI FILTERS 🛂

RESULTS BY YEAR



3 results



3 results

TEXT AVAILABILITY


Abstract

Free full text

Full text



Full text



Free PMC article.

PMID: 32209730

Share

Dy biginformatics analysis TET2 syntaggion was found positively correlated with tumor augustraces

Expression and prognosis analysis of *TET* family in acute myeloid leukemia

Tingjuan Zhang^{1,2,3,4,*}, Yangli Zhao^{2,5,*}, Yangjing Zhao⁶, Jingdong Zhou^{1,2,3,4}

- ¹ Department of Hematology, Affiliated People's Hospital of Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China
- ² Zhenjiang Medical School, Nanjing Medical University, Zhenjiang, Jiangsu, People's Republic of China
- ³ Zhenjiang Clinical Research Center of Hematology, Zhenjiang, Jiangsu, People's Republic of China
- ⁴ The Key Lab of Precision Diagnosis and Treatment in Hematologic Malignancies of Zhenjiang, Zhenjiang, Jiangsu, People's Republic of China
- ⁵ Department of Bioinformatics, School of Biomedical Engineering and Informatics, Nanjing Medical University, Nanjing, Jiangsu, People's Republic of China
- ⁶ Department of Immunology, Key Laboratory of Medical Science and Laboratory Medicine of Jiangsu Province, School of Medicine, Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China
- * Equal contribution

Received: December 30, 2019 Accepted: March 4, 2020 Published: March 25, 2020

https://doi.org/10.18632/aging.102928 How to Cite

Copyright © 2020 Zhang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

TET family members (TETs) encode proteins that represent crucial factors in the active DNA demethylation pathway. Evidence has proved that TET2 mutation is associated with leukemogenesis, drug response, and prognosis in acute myeloid leukemia (AML). However, few studies revealed the TETs expression and its clinical significance in AML. We conducted a detailed expression and prognosis analysis of TETs expression in human AML cell lines and patients by using public databases. We observed that TETs expression especially TET2 and TET3 was closely associated with AML among various human cancers. TET1 expression was significantly reduced in AML patients, whereas TET2 and TET3 expression was significantly increased. Kaplan-Meier analysis showed that only TET3 expression was associated with overall survival (OS) and disease-free survival (DFS) among both total AML as well as non-M3 AML, and was confirmed by another independent cohort. Moreover, Cox regression analysis revealed that TET3 expression may act as an independent prognostic factor for OS and DFS in total AML. Interestingly, patients that received hematopoietic stem cell transplantation (HSCT) did not show significantly longer OS and DFS than those who did not receive HSCT in TET3 high-expressed



Expression and prognosis analysis of *TET* family in acute myeloid leukemia

Tingjuan Zhang^{1,2,3,4,*}, Yangli Zhao^{2,5,*}, Yangjing Zhao⁶, Jingdong Zhou^{1,2,3,4}

- ¹ Department of Hematology, Affiliated People's Hospital of Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China
- ² Zhenjiang Medical School, Nanjing Medical University, Zhenjiang, Jiangsu, People's Republic of China
- ³ Zhenjiang Clinical Research Center of Hematology, Zhenjiang, Jiangsu, People's Republic of China
- ⁴ The Key Lab of Precision Diagnosis and Treatment in Hematologic Malignancies of Zhenjiang, Zhenjiang, Jiangsu, People's Republic of China
- ⁵ Department of Bioinformatics, School of Biomedical Engineering and Informatics, Nanjing Medical University, Nanjing, Jiangsu, People's Republic of China
- ⁶ Department of Immunology, Key Laboratory of Medical Science and Laboratory Medicine of Jiangsu Province, School of Medicine, Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China

Received: December 30, 2019 Accepted: March 4, 2020 Published: March 25, 2020

https://doi.org/10.18632/aging.102928 How to Cite

Copyright © 2020 Zhang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

TET family members (TETs) encode proteins that represent crucial factors in the active DNA demethylation pathway. Evidence has proved that TET2 mutation is associated with leukemogenesis, drug response, and prognosis in acute myeloid leukemia (AML). However, few studies revealed the TETs expression and its clinical significance in AML. We conducted a detailed expression and prognosis analysis of TETs expression in human AML cell lines and patients by using public databases. We observed that TETs expression especially TET2 and TET3 was closely associated with AML among various human cancers. TET1 expression was significantly reduced in AML patients, whereas TET2 and TET3 expression was significantly increased. Kaplan-Meier analysis showed that only TET3 expression was associated with overall survival (OS) and disease-free survival (DFS) among both total AML as well as non-M3 AML, and was confirmed by another independent cohort. Moreover, Cox regression analysis revealed that TET3 expression may act as an independent prognostic factor for OS and DFS in total AML. Interestingly, patients that received hematopoietic stem cell transplantation (HSCT) did not show significantly longer OS and DFS than those who did not receive HSCT in TET3 high-expressed

Does not talk about aging.

^{*} Equal contribution

Expression and prognosis analysis of *TET* family in acute myeloid leukemia

Tingjuan Zhang^{1,2,3,4,*}, Yangli Zhao^{2,5,*}, Yangjing Zhao⁶, Jingdong Zhou^{1,2,3,4}

- ¹ Department of Hematology, Affiliated People's Hospital of Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China
- ² Zhenjiang Medical School, Nanjing Medical University, Zhenjiang, Jiangsu, People's Republic of China
- ³ Zhenjiang Clinical Research Center of Hematology, Zhenjiang, Jiangsu, People's Republic of China
- ⁴ The Key Lab of Precision Diagnosis and Treatment in Hematologic Malignancies of Zhenjiang, Zhenjiang, Jiangsu, People's Republic of China
- ⁵ Department of Bioinformatics, School of Biomedical Engineering and Informatics, Nanjing Medical University, Nanjing, Jiangsu, People's Republic of China
- ⁶ Department of Immunology, Key Laboratory of Medical Science and Laboratory Medicine of Jiangsu Province, School of Medicine, Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China

Received: December 30, 2019 Accepted: March 4, 2020 Published: March 25, 2020

https://doi.org/10.18632/aging.102928 How to Cite

Copyright © 2020 Zhang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

TET family members (TETs) encode proteins that represent crucial factors in the active DNA demethylation pathway. Evidence has proved that TET2 mutation is associated with leukemogenesis, drug response, and prognosis in acute myeloid leukemia (AML). However, few studies revealed the TETs expression and its clinical significance in AML. We conducted a detailed expression and prognosis analysis of TETs expression in human AML cell lines and patients by using public databases. We observed that TETs expression especially TET2 and TET3 was closely associated with AML among various human cancers. TET1 expression was significantly reduced in AML patients, whereas TET2 and TET3 expression was significantly increased. Kaplan-Meier analysis showed that only TET3 expression was associated with overall survival (OS) and disease-free survival (DFS) among both total AML as well as non-M3 AML, and was confirmed by another independent cohort. Moreover, Cox regression analysis revealed that TET3 expression may act as an independent prognostic factor for OS and DFS in total AML. Interestingly, patients that received hematopoietic stem cell transplantation (HSCT) did not show significantly longer OS and DFS than those who did not receive HSCT in TET3 high-expressed

Does not talk about aging.

Does not talk about lifespan.

^{*} Equal contribution

Expression and prognosis analysis of *TET* family in acute myeloid leukemia

Tingjuan Zhang^{1,2,3,4,*}, Yangli Zhao^{2,5,*}, Yangjing Zhao⁶, Jingdong Zhou^{1,2,3,4}

- ¹ Department of Hematology, Affiliated People's Hospital of Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China
- ² Zhenjiang Medical School, Nanjing Medical University, Zhenjiang, Jiangsu, People's Republic of China
- ³ Zhenjiang Clinical Research Center of Hematology, Zhenjiang, Jiangsu, People's Republic of China
- ⁴ The Key Lab of Precision Diagnosis and Treatment in Hematologic Malignancies of Zhenjiang, Zhenjiang, Jiangsu, People's Republic of China
- ⁵ Department of Bioinformatics, School of Biomedical Engineering and Informatics, Nanjing Medical University, Nanjing, Jiangsu, People's Republic of China
- ⁶ Department of Immunology, Key Laboratory of Medical Science and Laboratory Medicine of Jiangsu Province, School of Medicine, Jiangsu University, Zhenjiang, Jiangsu, People's Republic of China

Received: December 30, 2019 Accepted: March 4, 2020 Published: March 25, 2020

https://doi.org/10.18632/aging.102928 How to Cite

Copyright © 2020 Zhang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

TET family members (TETs) encode proteins that represent crucial factors in the active DNA demethylation pathway. Evidence has proved that TET2 mutation is associated with leukemogenesis, drug response, and prognosis in acute myeloid leukemia (AML). However, few studies revealed the TETs expression and its clinical significance in AML. We conducted a detailed expression and prognosis analysis of TETs expression in human AML cell lines and patients by using public databases. We observed that TETs expression especially TET2 and TET3 was closely associated with AML among various human cancers. TET1 expression was significantly reduced in AML patients, whereas TET2 and TET3 expression was significantly increased. Kaplan-Meier analysis showed that only TET3 expression was associated with overall survival (OS) and disease-free survival (DFS) among both total AML as well as non-M3 AML, and was confirmed by another independent cohort. Moreover, Cox regression analysis revealed that TET3 expression may act as an independent prognostic factor for OS and DFS in total AML. Interestingly, patients that received hematopoietic stem cell transplantation (HSCT) did not show significantly longer OS and DFS than those who did not receive HSCT in TET3 high-expressed

Does not talk about aging.

Does not talk about lifespan.

Does not talk about longevity.

^{*} Equal contribution

Downregulation of LINC00665 suppresses the progression of lung adenocarcinoma via regulating miR-181c-5p/ZIC2 axis

Wei Wei¹, Xiaoliang Zhao¹, Jiang Liu², Zhenfa Zhang¹

¹ Department of Lung Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

² Department of Molecule Imaging and Nuclear Medicine in Diagnosis and Treatment, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

Received: March 4, 2021 Accepted: May 31, 2021 Published: July 7, 2021

https://doi.org/10.18632/aging.203240

How to Cite

Copyright: © 2021 Wei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Long non-coding RNA (IncRNA) LINC00665 was demonstrated to be upregulated in lung adenocarcinoma (LUAD) and target miR-181c-5p. ZIC2, which is upregulated in LUAD, serves as a putative target of miR-181c-5p. In this study, we aimed to reveal whether LINC00665 regulates miR-181c-5p/ZIC2 axis to promote LUAD progression. The results showed that LINC00665, HOXA1, ZIC2, and HOXA11 levels were increased in LUAD tissues, while miR-181c-5p level was decreased when compared to the adjacent normal tissues. High expression levels of LINC00665, ZIC2, HOXA1 and HOXA11, and low expression of miR-181c-5p were closely linked to poor prognosis of LUAD patients. Knockdown of LINC00665 induced obvious inhibitions in cell viability, clone formation, invasion and tumorigenesis in LUAD cells, whereas miR-181c-5p downregulation significantly neutralized these effects. In addition, downregulation of ZIC2 obviously reversed the enhancements of cell viability, clone formation, invasion and tumorigenesis induced by miR-181c-5p knockdown. In summary, the present study reveals that silencing of LINC00665 suppresses LUAD progression through targeting miR-181c-5p/ZIC2 axis.



Downregulation of LINC00665 suppresses the progression of lung adenocarcinoma via regulating miR-181c-5p/ZIC2 axis

Wei Wei¹, Xiaoliang Zhao¹, Jiang Liu², Zhenfa Zhang¹

¹ Department of Lung Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

² Department of Molecule Imaging and Nuclear Medicine in Diagnosis and Treatment, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

Received: March 4, 2021 Accepted: May 31, 2021 Published: July 7, 2021

https://doi.org/10.18632/aging.203240

How to Cite

Copyright: © 2021 Wei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Long non-coding RNA (IncRNA) LINC00665 was demonstrated to be upregulated in lung adenocarcinoma (LUAD) and target miR-181c-5p. ZIC2, which is upregulated in LUAD, serves as a putative target of miR-181c-5p. In this study, we aimed to reveal whether LINC00665 regulates miR-181c-5p/ZIC2 axis to promote LUAD progression. The results showed that LINC00665, HOXA1, ZIC2, and HOXA11 levels were increased in LUAD tissues, while miR-181c-5p level was decreased when compared to the adjacent normal tissues. High expression levels of LINC00665, ZIC2, HOXA1 and HOXA11, and low expression of miR-181c-5p were closely linked to poor prognosis of LUAD patients. Knockdown of LINC00665 induced obvious inhibitions in cell viability, clone formation, invasion and tumorigenesis in LUAD cells, whereas miR-181c-5p downregulation significantly neutralized these effects. In addition, downregulation of ZIC2 obviously reversed the enhancements of cell viability, clone formation, invasion and tumorigenesis induced by miR-181c-5p knockdown. In summary, the present study reveals that silencing of LINC00665 suppresses LUAD progression through targeting miR-181c-5p/ZIC2 axis.

Does not talk about aging.

Downregulation of LINC00665 suppresses the progression of lung adenocarcinoma via regulating miR-181c-5p/ZIC2 axis

Wei Wei¹, Xiaoliang Zhao¹, Jiang Liu², Zhenfa Zhang¹

¹ Department of Lung Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

² Department of Molecule Imaging and Nuclear Medicine in Diagnosis and Treatment, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

Received: March 4, 2021 Accepted: May 31, 2021 Published: July 7, 2021

https://doi.org/10.18632/aging.203240

How to Cite

Copyright: © 2021 Wei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Long non-coding RNA (IncRNA) LINC00665 was demonstrated to be upregulated in lung adenocarcinoma (LUAD) and target miR-181c-5p. ZIC2, which is upregulated in LUAD, serves as a putative target of miR-181c-5p. In this study, we aimed to reveal whether LINC00665 regulates miR-181c-5p/ZIC2 axis to promote LUAD progression. The results showed that LINC00665, HOXA1, ZIC2, and HOXA11 levels were increased in LUAD tissues, while miR-181c-5p level was decreased when compared to the adjacent normal tissues. High expression levels of LINC00665, ZIC2, HOXA1 and HOXA11, and low expression of miR-181c-5p were closely linked to poor prognosis of LUAD patients. Knockdown of LINC00665 induced obvious inhibitions in cell viability, clone formation, invasion and tumorigenesis in LUAD cells, whereas miR-181c-5p downregulation significantly neutralized these effects. In addition, downregulation of ZIC2 obviously reversed the enhancements of cell viability, clone formation, invasion and tumorigenesis induced by miR-181c-5p knockdown. In summary, the present study reveals that silencing of LINC00665 suppresses LUAD progression through targeting miR-181c-5p/ZIC2 axis.

Does not talk about aging.

Does not talk about lifespan.

Downregulation of LINC00665 suppresses the progression of lung adenocarcinoma via regulating miR-181c-5p/ZIC2 axis

Wei Wei¹, Xiaoliang Zhao¹, Jiang Liu², Zhenfa Zhang¹

¹ Department of Lung Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

² Department of Molecule Imaging and Nuclear Medicine in Diagnosis and Treatment, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

Received: March 4, 2021 Accepted: May 31, 2021 Published: July 7, 2021

https://doi.org/10.18632/aging.203240

How to Cite

Copyright: © 2021 Wei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Long non-coding RNA (IncRNA) LINC00665 was demonstrated to be upregulated in lung adenocarcinoma (LUAD) and target miR-181c-5p. ZIC2, which is upregulated in LUAD, serves as a putative target of miR-181c-5p. In this study, we aimed to reveal whether LINC00665 regulates miR-181c-5p/ZIC2 axis to promote LUAD progression. The results showed that LINC00665, HOXA1, ZIC2, and HOXA11 levels were increased in LUAD tissues, while miR-181c-5p level was decreased when compared to the adjacent normal tissues. High expression levels of LINC00665, ZIC2, HOXA1 and HOXA11, and low expression of miR-181c-5p were closely linked to poor prognosis of LUAD patients. Knockdown of LINC00665 induced obvious inhibitions in cell viability, clone formation, invasion and tumorigenesis in LUAD cells, whereas miR-181c-5p downregulation significantly neutralized these effects. In addition, downregulation of ZIC2 obviously reversed the enhancements of cell viability, clone formation, invasion and tumorigenesis induced by miR-181c-5p knockdown. In summary, the present study reveals that silencing of LINC00665 suppresses LUAD progression through targeting miR-181c-5p/ZIC2 axis.

Does not talk about aging.

Does not talk about lifespan.

Does not talk about longevity.

Downregulation of LINC00665 suppresses the progression of lung adenocarcinoma via regulating miR-181c-5p/ZIC2 axis

Wei Wei¹, Xiaoliang Zhao¹, Jiang Liu², Zhenfa Zhang¹

¹ Department of Lung Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

² Department of Molecule Imaging and Nuclear Medicine in Diagnosis and Treatment, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

Received: March 4, 2021 Ac

Accepted: May 31, 2021

Published: July 7, 2021

https://doi.org/10.18632/aging.203240

How to Cite

Copyright: © 2021 Wei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Long non-coding RNA (IncRNA) LINC00665 was demonstrated to be upregulated in lung adenocarcinoma (LUAD) and target miR-181c-5p. ZIC2, which is upregulated in LUAD, serves as a putative target of miR-181c-5p. In this study, we aimed to reveal whether LINC00665 regulates miR-181c-5p/ZIC2 axis to promote LUAD progression. The results showed that LINC00665, HOXA1, ZIC2, and HOXA11 levels were increased in LUAD tissues, while miR-181c-5p level was decreased when compared to the adjacent normal tissues. High expression levels of LINC00665, ZIC2, HOXA1 and HOXA11, and low expression of miR-181c-5p were closely linked to poor prognosis of LUAD patients. Knockdown of LINC00665 induced obvious inhibitions in cell viability, clone formation, invasion and tumorigenesis in LUAD cells, whereas miR-181c-5p downregulation significantly neutralized these effects. In addition, downregulation of ZIC2 obviously reversed the enhancements of cell viability, clone formation, invasion and tumorigenesis induced by miR-181c-5p knockdown. In summary, the present study reveals that silencing of LINC00665 suppresses LUAD progression through targeting miR-181c-5p/ZIC2 axis.

Does not talk about aging.

Does not talk about lifespan.

Does not talk about longevity.

Both appear in journal Aging-US.

Research Paper Volume 13, Issue 1

Volume 13, Issue 13 pp 17499—17515

Downregulation of LINC00665 suppresses the progression of lung adenocarcinoma via regulating miR-181c-5p/ZIC2 axis

Wei Wei¹, Xiaoliang Zhao¹, Jiang Liu², Zhenfa Zhang¹

¹ Department of Lung Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

² Department of Molecule Imaging and Nuclear Medicine in Diagnosis and Treatment, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin Lung Cancer Center, Tianjin 300060, China

Received: March 4, 2021

Accepted: May 31, 2021

Published: July 7, 2021

https://doi.org/10.18632/aging.203240

How to Cite

Copyright: © 2021 Wei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Long non-coding RNA (IncRNA) LINC00665 was demonstrated to be upregulated in lung adenocarcinoma (LUAD) and target miR-181c-5p. ZIC2, which is upregulated in LUAD, serves as a putative target of miR-181c-5p. In this study, we aimed to reveal whether LINC00665 regulates miR-181c-5p/ZIC2 axis to promote LUAD progression. The results showed that LINC00665, HOXA1, ZIC2, and HOXA11 levels were increased in LUAD tissues, while miR-181c-5p level was decreased when compared to the adjacent normal tissues. High expression levels of LINC00665, ZIC2, HOXA1 and HOXA11, and low expression of miR-181c-5p were closely linked to poor prognosis of LUAD patients. Knockdown of LINC00665 induced obvious inhibitions in cell viability, clone formation, invasion and tumorigenesis in LUAD cells, whereas miR-181c-5p downregulation significantly neutralized these effects. In addition, downregulation of ZIC2 obviously reversed the enhancements of cell viability, clone formation, invasion and tumorigenesis induced by miR-181c-5p knockdown. In summary, the present study reveals that silencing of LINC00665 suppresses LUAD progression through targeting miR-181c-5p/ZIC2 axis.

Does not talk about aging.


Does not talk about lifespan.

Does not talk about longevity.


Both appear in journal Aging-US.


Both talk about cancer.

Aging-US has been growing rapidly.




Aging-US has been growing rapidly.







—— Aging-US



Aging-US — Other geroscience journals






Is Aging-US close to research paper mills?

Is Aging-US close to research paper mills?

compile 56 lists of suspected paper mill products




n=25,420 publications

Ranking journals by citations toward suspected paper mill products retrieves Aging-US.

Journal	Citations
OncoTargets and therapy	254
International journal of molecular sciences	252
Cancer management and research	206
Journal of cellular and molecular medicine	183
Aging-US	170
Biomedicine & pharmacotherapy	164
Life Sciences	164
Oncology letters	162
Cancers	160

Ranking journals by citations toward suspected paper mill products retrieves Aging-US.




Journal	Citations
OncoTargets and therapy	254
International journal of molecular sciences	252
Cancer management and research	206
Journal of cellular and molecular medicine	183
Aging-US	170
Biomedicine & pharmacotherapy	164
Life Sciences	164
Oncology letters	162
Cancers	160

The National Science Library of the Chinese Academy of Sciences placed Aging-US on their journal warning list.

December 31st, 2021



Smut Clyde observed that formally independent publications in Aging-US share the same ruler.



https://forbetterscience.com/2021/05/17/the-ruler-of-the-aging-papermill/

Aging-US has been extraordinarily supportive of exciting directions within geroscience.

Aging-US has been extraordinarily supportive of exciting directions within geroscience.

Research Paper Volume 10, Issue 4

pp 573—591



An epigenetic biomarker of aging for lifespan and healthspan

Morgan E. Levine¹, Ake T. Lu¹, Austin Quach¹, Brian H. Chen², Themistocles L. Assimes³, Stefania Bandinelli⁴, Lifang Hou⁵, Andrea A. Baccarelli⁶, James D. Stewart⁷, Yun Li⁸, Eric A. Whitsel^{7,9}, James G Wilson¹⁰, Alex P Reiner¹¹, Abraham Aviv¹², Kurt Lohman¹², Yongmei Liu¹⁴, Luigi Ferrucci^{2, *}, Steve Horvath^{1,15, *}

Received: March 20, 2018 Accepted: April 8, 2018

Published: April 18, 2018

published some of the most innovative research

Aging-US has been extraordinarily supportive of exciting directions within geroscience.

Research Paper

Volume 10, Issue 4

pp 573-591



An epigenetic biomarker of aging for lifespan and healthspan

Morgan E. Levine¹, Ake T. Lu¹, Austin Quach¹, Brian H. Chen², Themistocles L. Assimes³, Stefania Bandinelli⁴, Lifang Hou⁵, Andrea A. Baccarelli⁶, James D. Stewart⁷, Yun Li⁸, Eric A. Whitsel^{7,9}, James G Wilson¹⁰, Alex P Reiner¹¹, Abraham Aviv¹², Kurt Lohman¹², Yongmei Liu¹⁴, Luigi Ferrucci^{2,*}, Steve Horvath^{1,15,*}

Received: March 20, 2018

Accepted: April 8, 2018

Published: April 18, 2018

published some of the most innovative research

























sponsors early stage conferences

Ambiguity

Ambiguity

Journal Hopping

Hero of the story: Reese Richardson https://reeserichardson.blog/




Hero of the story: Reese Richardson https://reeserichardson.blog/




On fall 2025 job market


OncoTarget is the biggest de-indexed biomedical journal.



OncoTarget is the biggest de-indexed biomedical journal.



OncoTarget is the biggest de-indexed biomedical journal.



Aging-US and OncoTarget are organizationally close.

Aging-US and OncoTarget are organizationally close.

From: Assistant Editor <u>@aging-us.com</u>>

Subject: Aging is associated with a systemic length-driven transcriptome imbalance

Date: December 6, 2022 at 08:25:45 CST

To: thomas.stoeger@northwestern.edu, a-misharin@northwestern.edu, s-buding@northwestern.edu, amaral@northwestern.edu

Cc: Mikhail Blagosklonny < blagosklonny@oncotarget.com >

Dear Authors,

I noticed your very interesting paper: Aging is associated with a systemic length-driven transcriptome imbalance.

I invite you to <u>submit</u> your paper to Aging (<u>https://aging.msubmit.net/cgi-bin/main.plex</u>), a multidisciplinary journal with an **Impact Factor of 5.955.**

Please submit your paper online. You may also submit it to editors@impactaging.com to accelerate consideration. If accepted, your paper will appear online and on MEDLINE/PubMed shortly after.

With warm regards,

Assistant Editor

On behalf of:

Prof. Mikhail Blagosklonny MD, PhD, Editor-in-Chief, www.aging-us.com

Aging-US and OncoTarget are organizationally close.

From: Assistant Editor @aging-us.com>

Subject: Aging is associated with a systemic length-driven transcriptome imbalance

Date: December 6, 2022 at 08:25:45 CST

To: thomas.stoeger@northwestern.edu, a-misharin@northweste

Cc: Mikhail Blagosklonny < blagosklonny@oncotarget.com >

Dear Authors,

Aging-US uses OncoTarget emails in their outreach.

I noticed your very interesting paper: Aging is associated with a systemic length-driven transcriptome imbalance.

I invite you to <u>submit</u> your paper to Aging (<u>https://aging.msubmit.net/cgi-bin/main.plex</u>), a multidisciplinary journal with an **Impact** Factor of 5.955.

Please submit your paper online. You may also submit it to editors@impactaging.com to accelerate consideration. If accepted, your paper will appear online and on MEDLINE/PubMed shortly after.


With warm regards,

Assistant Editor

On behalf of:


Prof. Mikhail Blagosklonny MD, PhD, Editor-in-Chief, www.aging-us.com

Many authors of Aging-US also published in OncoTarget.




—— Aging-US —— Other geroscience journals


Aging-US articles that cite suspected paper mill products disproportionally cite OncoTarget.




Aging-US articles that cite suspected paper mill products disproportionally cite OncoTarget.




Aging-US articles that cite suspected paper mill products disproportionally cite OncoTarget.






Is OncoTarget/Aging-US special?




Is OncoTarget/Aging-US special?



Our Services

Arda Conference is a global class interactive professional association for technical development and scientific knowledge platform where the international academics and professional community often meet and share their experience and wisdom in the forum that leads to its growth in addition to updates on cutting edge advancement on relevant research fields and technology.



Conferences and Meetings

Countless conferences take place around the world every day, with us, you stay updated on each one of them.



Journal Publications

We provide assistance with rapid journal publication with Scopus indexed journals, Web of Science, Peerreviewed, UGC and SCI/SCIE.



Article Writing Services

Whether you are a scholar or a student, we assist you with research article writing for UGC, Scopus, SCI, Web of Science Journals.



Research Consultancy

We've got your back even with Undergraduate, Post Graduate, Academic and Professional Research Activities.



PhD/Education Guidance

Dissertation and Thesis writing along with assistance to those pursuing Ph.D research through experienced mentor/guide.



Research Proposal Writing

We help our audience with choosing a good topic or presentation for their research proposal with original research proposal.

ARDA offers Journal Publication Services.

Journal Publication Services

ARDA offers research professionals who are desperate to have their research works known to the globe via publication in either Scopus Indexed Journals, UGC Care List Journals, Web of Science Journals, Peer-reviewed Journals / Google Scholar Journals or SCI/SCIE Indexed Journals, the chance to do so.

With its highly acclaimed publication assistance services, ARDA has helped research professionals from all across the world to fulfil their publication goals.

Scopus Indexed Journals

UGC Care List Journals

Web of Science Journals

Google Scholar journals | SCI / SCIE Indexed Journals















































Does de-indexing help?


Does de-indexing help?

Does de-indexing spread bad science?


Does de-indexing help?

Does de-indexing spread bad science?

What to do beyond de-indexing?







Spencer Hong



Jennifer Byrne



Luis Amaral

Everyone who flagged suspected paper mill products, especially:
Anna Abalkina, Elisabeth Bik,
Guillaume Cabanac, Leonid Schneider,
Retractionwatch, Smut Clyde, Tiger BB8

https://amaral.northwestern.edu

https://thomasstoegerlab.org